



A planar Schrödinger–Newton system with Trudinger–Moser critical growth

Zhisu Liu¹ · Vicențiu D. Rădulescu^{2,3,4,5} · Jianjun Zhang⁶

Received: 27 March 2022 / Accepted: 24 February 2023 / Published online: 20 March 2023
© The Author(s) 2023

Abstract

In this paper, we focus on the existence of positive solutions to the following planar Schrödinger–Newton system with general critical exponential growth

$$\begin{cases} -\Delta u + u + \phi u = f(u) & \text{in } \mathbb{R}^2, \\ \Delta \phi = u^2 & \text{in } \mathbb{R}^2, \end{cases}$$

where $f \in C^1(\mathbb{R}, \mathbb{R})$. We apply a variational approach developed in [36] to study the above problem in the Sobolev space $H^1(\mathbb{R}^2)$. The analysis developed in this paper also allows to investigate the relation between a Riesz-type of Schrödinger–Newton systems and a logarithmic-type of Schrödinger–Poisson systems. Furthermore, this approach can overcome some difficulties resulting from either the nonlocal term with sign-changing and unbounded logarithmic integral kernel, or the critical nonlinearity, or the lack of monotonicity of $\frac{f(t)}{t^3}$. We emphasize that it seems much difficult to use the variational framework developed in the existed literature to study the above problem.

Communicated by Andrea Mondino.

✉ Vicențiu D. Rădulescu
radulescu@inf.ucv.ro

Zhisu Liu
liuzhisu@cug.edu.cn

Jianjun Zhang
zhangjianjun09@tsinghua.org.cn

¹ Center for Mathematical Sciences/School of Mathematics and Physics, China University of Geosciences, Wuhan 430074, Hubei, People's Republic of China

² Faculty of Applied Mathematics, AGH University of Science and Technology, Al. Mickiewicza 30, 30-059 Kraków, Poland

³ Faculty of Electrical Engineering and Communication, Brno University of Technology, Technická 3058/10, 61600 Brno, Czech Republic

⁴ Department of Mathematics, University of Craiova, Street A.I. Cuza 13, 200585 Craiova, Romania

⁵ Simion Stoilow Institute of Mathematics of the Romanian Academy, P.O. Box 1-764, 014700 Bucharest, Romania

⁶ College of Mathematics and Statistics, Chongqing Jiaotong University, Chongqing 400074, People's Republic of China

Mathematics Subject Classification Primary 35Q55; Secondary 35B33 · 35J47 · 46E35

1 Introduction and results

1.1 Overview

Consider the following nonlinear Schrödinger–Newton system

$$\begin{cases} i\hbar \frac{\partial \psi}{\partial t} = \frac{\hbar^2}{2m} \Delta \psi + W(x)\psi + \lambda \phi \psi - f(\psi) & \text{in } \mathbb{R}^d \times \mathbb{R}, \\ \Delta \phi = |\psi|^2 & \text{in } \mathbb{R}^d \times \mathbb{R}, \end{cases} \tag{1.1}$$

where $\lambda \in \mathbb{R}$, i is the imaginary unit, \hbar is the Planck constant. For $d = 3$, $m > 0$ stands for the mass of the particle, $\psi : \mathbb{R}^3 \times [0, T] \rightarrow \mathbb{C}$ is a wave function, W is a real external potential and such a system often appears in quantum mechanics models and semiconductor theory (see [33]) and also arises, for example, as a model of the interaction of a charged particle with the electrostatic field (see [7]). It is well known that $\psi(x, t) = u(x)e^{-\frac{iEt}{\hbar}}$, $x \in \mathbb{R}^d$, $t \in \mathbb{R}$ is a standing wave solution of (1.1) if and only if $u : \mathbb{R}^d \rightarrow \mathbb{R}$ satisfies

$$\begin{cases} -\frac{\hbar^2}{2m} \Delta u + (W(x) - E)u + \lambda \phi u = f(u) & \text{in } \mathbb{R}^d, \\ \Delta \phi = u^2 & \text{in } \mathbb{R}^d. \end{cases} \tag{1.2}$$

The second equation in system (1.2) can be solved by

$$\phi(x) = \Gamma_d(x) * u^2(x) = \int_{\mathbb{R}^d} \Gamma_d(x - y)u^2(y)dy,$$

where Γ_d is the Newtonian kernel in dimension d , which is expressed by

$$\Gamma_d(x) = \begin{cases} \frac{1}{2\pi} \ln |x|, & d = 2, \\ \frac{1}{d(2-d)\omega_d} |x|^{2-d}, & d \geq 3. \end{cases}$$

Here ω_d is the volume of the unit d -ball. Under such a formal inversion of the second equation in (1.2), we obtain the following non-local equation

$$-\Delta u + V(x)u + \lambda(\Gamma_d * |u|)u = f(u) \quad \text{in } \mathbb{R}^d, \tag{1.3}$$

where $V = W - E$. The cases $\lambda > 0$ and $\lambda < 0$ denote respectively two very different physical situations(see [22]). In particular, when $\lambda > 0$, (1.3) stands for one attractive case of a Newton-Poisson coupling for gravitational mean-field models. When $\lambda < 0$, (1.3) represents one d -dimensional case of repulsive electrostatic forces. Problem (1.3) is variational formally, and its associated energy functional is given by

$$\begin{aligned} I_d(u) &= \frac{1}{2} \int_{\mathbb{R}^d} (|\nabla u|^2 + V(x)u^2) dx + \frac{\lambda}{4} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \Gamma_d(|x - y|) u^2(x)u^2(y) dx dy \\ &\quad - \int_{\mathbb{R}^d} F(u) dx. \end{aligned}$$

In the case $d = 3$, I_d is well defined and of C^1 class in $H^1(\mathbb{R}^d)$ when $V \in L^\infty(\mathbb{R}^d)$. In the literature, by exploring the variational methods and topological methods, the existence, nonexistence, multiplicity and concentration of solutions to (1.3) have been investigated when f and V satisfy various assumptions, see e.g. [5, 7, 26, 28, 34, 38, 41] and so on.

Throughout this paper, we assume $\frac{\lambda}{2\pi} = 1$, and consider the following Schrödinger–Newton equation

$$-\Delta u + V(x)u + (\ln(|\cdot|) * |u|^2)u = f(u) \text{ in } \mathbb{R}^2, \tag{1.4}$$

whose formal energy functional can be given by

$$I(u) = \frac{1}{2} \int_{\mathbb{R}^2} (|\nabla u|^2 + V(x)u^2) dx + \frac{1}{4} \int_{\mathbb{R}^4} \ln(|x - y|) u^2(x)u^2(y) dx dy - \int_{\mathbb{R}^2} F(u) dx.$$

Since $\tilde{\Gamma}(x) := \ln|x|$ is sign-changing and presents singularities at zero and infinity, compared with the higher dimensional case $d \geq 3$, the associated energy functional with (1.4) seems much more delicate. In particular, functional I is not well-defined on $H^1(\mathbb{R}^2)$ because of the appearance of the singular convolution term

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(|x - y|) u^2(x)u^2(y) dx dy,$$

which is not well defined for all $u \in H^1(\mathbb{R}^2)$. Therefore, the approaches dealing with higher dimensional cases seem difficult to be adapted to the case $d = 2$. So the rigorous study of the planar Schrödinger–Newton system had remained open for a long time. Recall that Choquard, Stubbe and Vuffray [19] proved the existence of a unique positive radially symmetric solution to (1.4) with $V(x) \equiv 1$ and $f(x, u) = 0$ by applying a shooting method. To consider problem (1.4) with $d = 2$ and $V(x) \equiv 1$, Stubbe [39] introduced the following weighted Sobolev space

$$X := \left\{ u \in H^1(\mathbb{R}^2) : \int_{\mathbb{R}^2} \ln(1 + |x|) |u(x)|^2 dx < +\infty \right\},$$

endowed with the norm

$$\|u\|_X^2 = \int_{\mathbb{R}^2} (|\nabla u|^2 + |u|^2) dx + \int_{\mathbb{R}^2} \ln(1 + |x|) |u(x)|^2 dx,$$

which yields that the associated energy functional is well-defined and continuously differentiable on the space X . More precisely, thanks to the Hardy–Littlewood–Sobolev inequality [29], for any $u \in X$,

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(|x - y|) u^2(x)u^2(y) dx dy$$

can be controlled by

$$\int_{\mathbb{R}^2} \ln(1 + |x|) u^2(x) dx.$$

Consequently, within the underlying space X above, Cingolani and Weth [20] studied problem (1.4) with $f(u) = |u|^{p-2}u$, $p \geq 4$ and obtained the existence and multiplicity of solutions. In studying planar Schrödinger–Newton systems in the underlying space X , one of main obstacles is that the norm $\|\cdot\|_X$ lacks translation invariance. This makes problems tough in verifying the compactness via the concentration-compactness principle. In [20, Lemma 2.1], it is shown that this difficulty can be overcome via a symmetric bilinear form

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(1 + |x - y|) u(x)v(y) dx dy.$$

In a similar fashion, a sequence of higher energy solutions was obtained in [20] for $p \geq 4$ in a periodic setting, where the corresponding energy functional is invariant under \mathbb{Z}^2 -translations. Later, Du and Weth [23] extended the above results to the case $p \in (2, 4)$. Under the above variational framework in [20, 39], Chen and Tang in [16] considered the planar Schrödinger–Newton system in the axially symmetric setting. By using Jeanjean’s monotonicity trick [27] and a Nehari–Pohozaev manifold argument, they proved that there exists at least a ground state solution to (1.4). For some other related works to the two dimensional case, see [6, 9, 13, 17, 18, 21, 42] and the references therein. In all the results mentioned above for the planar Schrödinger–Newton system, it is obvious that the weighted function space X plays a fundamental role in ensuring that the energy functional is well defined and continuously differentiable.

Different from the variational frameworks above, the authors in [36] introduce a novel variational approach to study problem (1.4) by considering a perturbation problem defined in $H^1(\mathbb{R}^2)$. We aim to use a variational approach established in [36] to problem (1.4) involving the critical exponential growth in the sense of Trudinger–Moser, see [37, 40]. We now recall a notion of criticality which is totally different from the Sobolev type.

(f_0) there exists $\theta_0 > 0$ such that

$$\lim_{|t| \rightarrow \infty} \frac{f(t)}{e^{\theta t^2}} = 0, \quad \forall \theta > \theta_0, \quad \lim_{|t| \rightarrow \infty} \frac{f(t)}{e^{\theta t^2}} = +\infty, \quad \forall \theta < \theta_0,$$

which was introduced by Adimurthi and Yadava [2] and see also de Figueiredo, Miyagaki and Ruf [24]. We stress that Alves and Figueiredo [4] investigated the existence of positive ground state solutions for (1.4) when $V(x) \equiv 1$ and f satisfies (f_0) and the following conditions:

- (f_1) $f \in C(\mathbb{R}, \mathbb{R})$ and $f(t) = o(t)$ as $t \rightarrow 0$.
- (f_2) $\frac{f(t)}{t^3}$ is increasing in $(0, \infty)$;
- (f_3) there exists $\mu > 2$ such that $0 < \mu F(t) \leq f(t)t$ for all $t > 0$, where $F(t) = \int_0^t F(s)ds$;
- (f_4) there exist constants $p > 4$ and $\lambda_0 > c_p$ for some positive constant c_p depending on p .

And later, Chen and Tang [17] studied the existence of nontrivial solutions to (1.4) when $f(u)$ is replaced by $f(x, u) \in C(\mathbb{R}^2 \times \mathbb{R}, \mathbb{R})$ which is required to satisfy the following conditions:

- (F_1) $f(x, t) = o(t)$ as $t \rightarrow 0$ uniformly for $x \in \mathbb{R}^2$; $f(x, t) = f(|x_1|, |x_2|, t) = f(|x_1|, |x_2|, t)$ for all $(x, t) \in \mathbb{R}^2 \times \mathbb{R}$.
- (F_2) $f(x, t)t > 0$ for all $(x, t) \in \mathbb{R}^2 \times (\mathbb{R} \setminus \{0\})$ and there exist $M_0 > 0$ and $t_0 > 0$ such that

$$F(x, t) \leq M_0 |f(x, t)|, \quad \forall x \in \mathbb{R}^2, |t| \geq t_0,$$

where $F(x, t) = \int_0^t F(x, s)ds$.

- (F_3) $\liminf_{t \rightarrow \infty} \frac{t^2 F(x, t)}{e^{\theta_0 t^2}} \geq \kappa > \frac{2}{\theta_0^2 \rho^2}$, where $\rho \in (0, 1/2)$ such that $\rho^2 \max_{|x| \leq \rho} V(x) \leq 1$.

- (F_4) For all $x \in \mathbb{R}^2$, the mapping $(0, \infty) \ni t \mapsto \frac{f(x, t) - V(x)t}{t^3}$ is non-decreasing.

1.2 Main result

Since we study the planar Schrödinger–Newton system with critical exponential nonlinearities in the sense of Trudinger–Moser, we first recall the 2D-Pohozaev–Trudinger–Moser

inequality, which was established by Cao [12], see also [1, 11, 14, 15]. This result is crucial in estimating the subcritical or critical nonlinearity of Trudinger–Moser type.

Lemma 1.1 [12] *If $\theta > 0$ and $u \in H^1(\mathbb{R}^2)$, then*

$$\int_{\mathbb{R}^2} (e^{\theta u^2} - 1) \, dx < \infty.$$

If, moreover, $u \in H^1(\mathbb{R}^2)$, $\|\nabla u\|_2^2 \leq 1$, $\|u\|_2^2 < M < \infty$ and $\theta < 4\pi$, then there exists a constant $C_{M,\theta}$ which depends only on M, θ such that

$$\int_{\mathbb{R}^2} (e^{\theta u^2} - 1) \, dx \leq C_{M,\theta}.$$

For this purpose, we make the following assumptions on the nonlinearity $f \in C^1(\mathbb{R}, \mathbb{R})$.

(f₅) $f(t)t > 0$ for all $t \in \mathbb{R} \setminus \{0\}$, and there exist $M_0 > 0$ and $t_0 > 0$ such that

$$F(t) \leq M_0|f(t)|, \quad \forall |t| \geq t_0,$$

where $F(t) = \int_0^t f(s) \, ds$.

(f₆) $\liminf_{t \rightarrow \infty} \frac{t^2 F(t)}{e^{\theta_0 t^2}} \geq \kappa > \frac{1}{4\sqrt{\theta_0 \pi} \rho^2}$ for $\rho \in (0, 1/2)$, where θ_0 is given in (f₀).

(f₇) The function $\frac{f(t)t - F(t)}{t^3}$ is nondecreasing in $(0, +\infty)$.

Remark 1.2 It follows from conditions (f₁), (f₅) and (f₇) that $0 < 3F(s) \leq f(s)s$ for $s > 0$.

Since we aim at finding positive solutions of equation (1.4), we always assume $f(s) \equiv 0$ for $s \leq 0$, throughout this paper.

Our main results states as follows.

Theorem 1.3 *Assume that hypotheses (f₀)–(f₁) and (f₅)–(f₇) hold. Then equation (1.4) with $V(x) \equiv 1$ has at least a positive solution $u \in H^1(\mathbb{R}^2)$ satisfying*

$$\left| \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| u^2(x) u^2(y) \, dx \, dy \right| < +\infty. \tag{1.5}$$

Remark 1.4 Observe from [4] that the monotonicity condition (f₂) is often used to guarantee the boundedness of the Palais–Smale sequence $\{u_n\}$. With the aid of (f₄), the authors in [4] established directly an upper estimate on the $H^1(\mathbb{R}^2)$ -norm of Palais–Smale sequence $\{u_n\}$. Then thanks to the Trudinger–Moser inequality, the compactness is recovered. However, as a global condition, (f₄) requires $f(t)$ to be super-cubic for all $t \geq 0$, which seems a little bit strict especially for $t > 0$ small. Observe that condition (f₄) does not reveal the essential features of the exponential growth given in (f₀). As mentioned in [17], there exist many model nonlinearities without satisfying (f₂) or (f₄) which are required in [4].

Observe that Chen and Tang in [17] obtained the existence of nontrivial solutions under (F₁)–(F₄) which are weaker than those in [4]. Moreover, the authors in [17] introduced conditions (F₂) and (F₃) to state an upper estimate for the minimax-level using the Moser type sequence, so that vanish does not occur for the Cerami sequence $\{u_n\}$. However, in order to prove that the weak limit function \bar{u} of Cerami sequence $\{u_n\}$ is a solution of system (1.4), one need to show directly

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x - y| u_n^2 \, dy u_n \phi \, dx = \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln(|x - y|) \bar{u}^2 \, dy \bar{u} \phi \, dx, \quad \forall \phi \in C_0^\infty(\mathbb{R}^2) \tag{1.6}$$

and

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} f(x, u_n)(u_n - \bar{u})dx = 0 \tag{1.7}$$

without (f_2) or (f_4) , which seems tough to establish in the weight space X , even if $u_n \rightarrow \bar{u}$ in $L^s(\mathbb{R}^2)$ for $s \in [2, +\infty)$. And so, (F_4) in [17] was introduced to guarantee that the associated energy functional can be studied in Nehari-type manifold, and then use some energy estimate method together with Fatou’s lemma to recover compactness in space X . We emphasize that (F_4) in [17] plays an essential role in proving the existence of nontrivial solution.

In the present paper, we also need (f_5) and (f_6) to establish a similar upper estimate as [17] by using the Moser type sequence. However, (f_7) is weaker than (F_4) , when we consider autonomous nonlinearity f . One can not restrict functional I on Nehari-type manifold to study directly, since (f_7) results in that I has no lower bound at Nehari-type manifold. It even seems difficult to find some suitable manifold in the weighted space X to use constraint variational approaches to obtain (1.6) and (1.7) under condition (f_7) .

Remark 1.5 Very recently, Albuquerque et al., [3] investigated the existence of solutions to the planar non-autonomous Schrödinger–Poisson system

$$\begin{cases} -\Delta u + V(|x|)u + \gamma\phi K(|x|)u = \lambda Q(|x|)f(u), & x \in \mathbb{R}^2, \\ \Delta\phi = K(|x|)u^2, & x \in \mathbb{R}^2, \end{cases} \tag{1.8}$$

where γ, λ are positive parameters, V, K, Q are continuous potentials, which can be unbounded or vanishing at infinity. By assuming that the nonlinearity $f(t)$ satisfies (f_0) , (f_2) and

(\tilde{f}_1) $f(s) = o(|s|^{\gamma-1})$ as $s \rightarrow 0$, where

$$\gamma := \max\{2, 2(2 + 2b - a)/(a + 2)\} = \begin{cases} 2 & \text{if } -2 < b \leq a, \\ 2(2 + 2b - a)/(a + 2) & \text{if } -2 < a < b, \end{cases}$$

(\tilde{f}_2) there exists $\theta > \max\{\gamma, 4\}$ such that $0 < \theta F(s) \leq f(s)s$ for all $s \geq 0$,

(\tilde{f}_3) there exists $q > \gamma$ such that $\liminf_{s \rightarrow 0^+} F(s)/s^q > 0$,

under a similar variational framework as that in [20], they derived the existence of a ground state solution to system (1.8) for λ large enough. Compared with [3], we use the different variational framework to weaken the conditions (\tilde{f}_2) and (\tilde{f}_3) .

1.3 Main difficulty and strategy

In the present paper, we employ the variational framework established in [36] to study problem (1.4) in the standard Sobolev space $H^1(\mathbb{R}^2)$ by variational methods. In order to overcome the difficulty that the sign-changing property of the Newtonian kernel $\Gamma_d(x) = \frac{1}{2\pi} \ln|x|$ leads to failure in setting the variational framework in $H^1(\mathbb{R}^2)$, as in [36], we modified equation (1.4) as follows

$$-\Delta u + u - \frac{1}{\alpha}(G_\alpha(|\cdot|) * u^2)u = f(u) \quad \text{in } \mathbb{R}^2, \tag{1.9}$$

where $\alpha \in (0, 1)$ is a parameter and

$$\lim_{\alpha \rightarrow 0^+} G_\alpha(x) := \lim_{\alpha \rightarrow 0^+} \frac{|x|^{-\alpha} - 1}{\alpha} = \ln|x|$$

for $x \in \mathbb{R}^2 \setminus \{0\}$. The corresponding energy functional to (1.9) is well defined in $H^1(\mathbb{R}^2)$ for fixed $\alpha \in (0, 1)$, which enables us to use minimax methods to study the existence of positive solutions for (1.9). By passing to the limit, a convergence argument within $H^1(\mathbb{R}^2)$ allows us to get positive solutions of the original problem (1.4).

In the limit process above as $\alpha \rightarrow 0^+$, the main difficulties are two-fold. Firstly, there is the lack of compactness due to the effect of critical exponential nonlinearity and the appearance of singularity at $\alpha = 0$. Secondly, the boundedness of the Palais–Smale sequences is not easy to get, since 4-Ambrosetti–Rabinowitz condition does not hold. Moreover, Jeanjean’s monotonicity trick [27] seems not to work at our problem, since the singularity at $\alpha = 0$ leads to failure at giving a uniform upper bound to the corresponding minimax value as $\alpha \rightarrow 0^+$.

In order to overcome these obstacles, in the proof of Theorem 1.3 we firstly adopt the perturbation introduced in [36](see also [34, 35]) to obtain the boundedness of the Palais–Smale sequences. Secondly, we need to use Moser type sequence together with some refined analysis to establish an upper estimate as a threshold to recover compactness locally. Thirdly, we use the concentration–compactness principle to establish a compactness splitting lemma of critical exponential version, and then to prove the modified equation (1.9) has a positive mountain pass solution u_α . Moreover, the mountain pass value c_α is uniformly bounded from below and above as $\alpha \rightarrow 0^+$. Lastly, it follows from the moving plane arguments that u_α is radially symmetric, and then one exponential decay of u_α at infinity can be obtained uniformly for $\alpha > 0$ small. Therefore, the Lebesgue dominated convergence theorem enables us to get the Frechet derivative of the corresponding energy functional is weakly sequence continuous and then get compactness.

Among other things our results will give the following findings and consequences:

- We use a variational approach (see also [36]) to study system (1.4) directly in the usual Sobolev space $H^1(\mathbb{R}^2)$, which is totally different from the one established in [20, 39]. Compared with solutions obtained in the weighted space X in the literature, we obtain solution u of system (1.4) in $H^1(\mathbb{R}^2)$ directly. Moreover, in our arguments we can find a relation between a Riesz-type of Schrödinger–Newton systems and a logarithmic-type of Schrödinger–Poisson systems.
- As mention in Remark 1.4, it seems tough to prove (1.6) and (1.7) directly in the weighted space X in our setting. That is to say, it seems difficult to use the variational approach established in [20] to prove Theorem 1.3. Therefore, this shows that the variational approach established in [36] can also be used to deal with some cases in which the variational approach [20] seems not easy to be adopted for us.

This paper is organized as follows. Some preliminaries are given in Section 2, and Section 3 is devoted to the existence of mountain pass type solutions to the modified equation. Then in Section 4, we complete the proof of Theorem 1.3.

2 Preliminary results

Let us fix some notations. The letter C will be repeatedly used to denote various positive constants, whose exact values may be irrelevant. Denote infinitely small quantities $o(1)$ and $o(\alpha)$ by $o(1) \rightarrow 0$ as $n \rightarrow \infty$ and $o(\alpha) \rightarrow 0$ as $\alpha \rightarrow 0^+$, respectively. For every $1 \leq s \leq +\infty$, we denote by $\|\cdot\|_s$ the usual norm of the Lebesgue space $L^s(\mathbb{R}^2)$. The function space

$$H^1(\mathbb{R}^2) := \{u \in L^2(\mathbb{R}^2) : |\nabla u| \in L^2(\mathbb{R}^2)\}$$

is the usual Sobolev space endowed with the norm

$$\|u\| := \left(\int_{\mathbb{R}^2} (|\nabla u|^2 + u^2) dx \right)^{\frac{1}{2}}.$$

In what follows, we recall the Hardy–Littlewood–Sobolev inequality (see [29]), which will be frequently used throughout this paper.

Lemma 2.1 (Hardy–Littlewood–Sobolev inequality [29]) *Let $s, r > 1$ and $\alpha \in (0, d)$ with $1/s + \alpha/d + 1/r = 2$, $f \in L^s(\mathbb{R}^d)$ and $h \in L^r(\mathbb{R}^d)$. There exists a sharp constant $C_{s,d,\alpha,r}$ independent of f, h , such that*

$$\int_{\mathbb{R}^d} \left[\frac{1}{|x|^\alpha} * f(x) \right] h(x) dx \leq C_{s,d,\alpha,r} \|f\|_s \|h\|_r.$$

If $r = s = \frac{2d}{2d-\alpha}$, then

$$C_{s,d,\alpha,r} = C_{d,\alpha} = \pi^{\alpha/2} \frac{\Gamma\left(\frac{d}{2} - \frac{\alpha}{2}\right)}{\Gamma\left(d - \frac{\alpha}{2}\right)} \left\{ \frac{\Gamma\left(\frac{d}{2}\right)}{\Gamma(d)} \right\}^{-1 + \frac{\alpha}{d}},$$

and if $d = 2, \alpha \in (0, 1]$, then $C_{2,\alpha} \leq 2\sqrt{\pi}$.

Lemma 2.2 (Moser-Trudinger inequality [1, 12]) *For any $\beta \in (0, 4\pi)$ there exists $C = C_\beta > 0$ such that for every $u \in H^1(\mathbb{R}^2)$ satisfying $\int_{\mathbb{R}^2} |\nabla u|^2 \leq 1$, one has*

$$\int_{\mathbb{R}^2} \min\{1, u^2\} e^{\beta|u|^2} \leq C_\beta \int_{\mathbb{R}^2} |u|^2.$$

We recall the following elementary lemma which is of use in doing energy estimate.

Lemma 2.3 [44, Lemma 2.1] *For any $\beta \in (0, \infty)$, there exists $C_\beta > 0$ such that*

$$\frac{s^{-\alpha} - 1}{\alpha} \leq C_\beta s^{-\beta}, \quad s > 0$$

holds for all $\alpha \in (0, \beta)$.

3 The modified problem

Since the fact that I is not well defined on $H^1(\mathbb{R}^2)$, we use the perturbation technique (see [36]) to overcome this difficulty by modifying Schrödinger–Newton systems. We state the following modified problem

$$-\Delta u + u - \frac{1}{\alpha} (G_\alpha(x) * u^2)u = f(u), \quad x \in \mathbb{R}^2, \tag{3.1}$$

where $\alpha \in (0, 1)$ is a parameter and $G_\alpha(x) = \frac{|x|^{-\alpha-1}}{\alpha}, x \neq 0$. Its associated functional is

$$I_\alpha(u) = \frac{1}{2} \|u\|^2 - \frac{1}{4} \int_{\mathbb{R}^2} (G_\alpha(x) * u^2)u^2 dx - \int_{\mathbb{R}^2} F(u) dx, \quad u \in H^1(\mathbb{R}^2).$$

By virtue of the definition of G_α , it follows from the Hardy-Littlewood-Sobolev inequality that for any given α , the perturbation functional I_α is well-defined on $H^1(\mathbb{R}^2)$, of C^1 -class and

$$I'_\alpha(u)v = \int_{\mathbb{R}^2} (\nabla u \nabla v + uv) dx - \int_{\mathbb{R}^2} (G_\alpha(x) * u^2)uv dx - \int_{\mathbb{R}^2} f(u)v dx$$

for $u, v \in H^1(\mathbb{R}^2)$. Since the conditions of Theorem 1.3 do not include the well-known 4-Ambrosetti–Rabinowitz condition, the boundedness of the Palais–Smale sequence is not easy to get. In order to overcome this difficulty, we add another perturbation technique developed in [34, 35] to equation (3.1). We now give more details to describe such a technique. Set

$$\lambda \in \left(0, \min \left\{ 1, \sqrt[4]{\frac{\theta_0}{5\pi}} \right\} \right), \quad r \in (4, +\infty).$$

Let us consider the following modified problem

$$-\Delta u + u - (G_\alpha(x) * u^2)u + \lambda \left(\int_{\mathbb{R}^2} u^2 dx \right)^{\frac{1}{4}} u = f(u) + \lambda |u|^{r-2}u, \quad u \in H^1(\mathbb{R}^2). \quad (3.2)$$

The associated functional with (3.2) is given by

$$I_{\alpha,\lambda}(u) = \frac{1}{2} \|u\|^2 - \frac{1}{4} \int_{\mathbb{R}^2} (G_\alpha(x) * u^2)u^2 dx + \frac{2\lambda}{5} \|u\|_{\frac{5}{2}}^{\frac{5}{2}} - \int_{\mathbb{R}^2} F(u) dx - \frac{\lambda}{r} \|u\|_r^r.$$

It is not hard for fixed $\alpha > 0$ to show that functional $I_{\alpha,\lambda}$ is well-defined on $H^1(\mathbb{R}^2)$, of C^1 -class and

$$I'_{\alpha,\lambda}(u)v = I'_\alpha(u)v + \lambda \|u\|_{\frac{5}{2}}^{\frac{1}{2}} \int_{\mathbb{R}^2} uv dx - \lambda \int_{\mathbb{R}^2} |u|^{r-2}uv dx$$

for $u, v \in H^1(\mathbb{R}^2)$. For any critical point $u \in H^1(\mathbb{R}^2)$ of $I_{\alpha,\lambda}$, the following Pohozaev identity holds

$$P_{\alpha,\lambda}(u) := \|u\|_2^2 + \frac{1}{\alpha} \|u\|_2^4 - \frac{4-\alpha}{4\alpha} Q(u) + \lambda \|u\|_{\frac{5}{2}}^{\frac{5}{2}} - 2 \int_{\mathbb{R}^2} \left(F(u) + \frac{\lambda}{r} |u|^r \right) dx = 0, \quad (3.3)$$

where $Q(u) = \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u^2 \right) u^2 dx$.

Lemma 3.1 *Suppose (f_0) – (f_1) and (f_5) – (f_7) hold, then*

- (i) *there exist $\rho, \delta_0 > 0$ (independent of α, λ) such that $I_{\alpha,\lambda}|_{S_\rho}(u) \geq \delta_0$ for every $u \in S_\rho = \{u \in H^1(\mathbb{R}^2) : \|u\| = \rho\}$;*
- (ii) *there is $e \in H^1(\mathbb{R}^2)$ (independent of α, λ) with $\|e\| > \rho$ such that $I_{\alpha,\lambda}(e) < 0$.*

Proof (i) From (f_0) – (f_1) , we have for any $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that

$$|F(t)| \leq \varepsilon |t|^2 + C_\varepsilon |t|^3 (e^{\theta_0 t^2} - 1). \quad (3.4)$$

Take $u \in H^1(\mathbb{R}^2)$ and $\|u\|^2 < 2\pi/\theta_0$. Obviously, $\int_{\mathbb{R}^2} |\nabla u|^2 dx < 2\pi/\theta_0$. So by Lemma 1.1, one has

$$\begin{aligned} \int_{\mathbb{R}^2} F(u) dx &\leq \varepsilon \int_{\mathbb{R}^2} |u|^2 dx + C_\varepsilon \int_{\mathbb{R}^2} |u|^3 (e^{\theta_0 u^2} - 1) dx \\ &\leq \varepsilon \|u\|^2 + C_\varepsilon \left[\int_{\mathbb{R}^2} (e^{2\theta_0 u^2} - 1) \right]^{1/2} \|u\|_6^3 \\ &\leq \varepsilon \|u\|^2 + C_\varepsilon \|u\|^3. \end{aligned} \quad (3.5)$$

It follows from (3.5), Lemma 2.3, and Hardy–Littlewood–Sobolev’s inequality that

$$\begin{aligned}
 I_{\alpha,\lambda}(u) &= \frac{1}{2}\|u\|^2 - \frac{1}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|x-y|^{-\alpha} - 1}{\alpha} u^2(y)u^2(x)dydx \\
 &\quad - \int_{\mathbb{R}^2} F(u)dx + \frac{2\lambda}{5}\|u\|_{\frac{5}{2}}^{\frac{5}{2}} - \frac{\lambda}{r}\|u\|_r^r \\
 &\geq \frac{1-2\varepsilon}{2}\|u\|^2 - \frac{1}{4} \int_{|x-y|\leq 1} \frac{|x-y|^{-\alpha} - 1}{\alpha} u^2(y)u^2(x)dydx - C_\varepsilon\|u\|^3 - \frac{\lambda}{r}\|u\|_r^r \\
 &\geq \frac{1-2\varepsilon}{2}\|u\|^2 - \frac{C}{4}\|u\|^4 - C_\varepsilon\|u\|^3 - C\frac{1}{r}\|u\|_r^r.
 \end{aligned}
 \tag{3.6}$$

Letting $\varepsilon \in (0, \frac{1}{2})$ and $\|u\| = \rho > 0$ small enough, it is clear that there exists $\delta_0 > 0$ such that $I_\alpha(u) \geq \delta_0$ for every $u \in S_\rho$.

(ii) The proof is very similar to that of Lemma 3.3 in [36]. For the reader’s convenience, we give the details. Take $e_0 \in C_0^\infty(\mathbb{R}^2)$ such that $e_0(x) \equiv 1$ for $x \in B_{\frac{1}{8}}(0)$, $e_0(x) \equiv 0$ for $x \in \mathbb{R}^2 \setminus B_{\frac{1}{4}}(0)$ and $|\nabla e_0(x)| \leq C$. Note that

$$\frac{s^{-\alpha} - 1}{\alpha} \geq \ln \frac{1}{s}, \quad \text{for } s \in (0, 1].$$

It then follows from the definition of $I_{\alpha,\lambda}$ that for $t > 0$,

$$\begin{aligned}
 I_{\alpha,\lambda}(te_0) &\leq \frac{t^2}{2}\|e_0\|^2 + \frac{2t^{5/2}}{5}\|u\|_{\frac{5}{2}}^{\frac{5}{2}} - \frac{t^4}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|x-y|^{-\alpha} - 1}{\alpha} e_0^2(y)e_0^2(x)dydx - \int_{\mathbb{R}^2} F(te_0)dx \\
 &\leq \frac{t^2}{2}\|e_0\|^2 - \frac{t^4}{4} \int_{|x|\leq \frac{1}{4}} \int_{|y|\leq \frac{1}{4}} \frac{|x-y|^{-\alpha} - 1}{\alpha} e_0^2(y)e_0^2(x)dydx \\
 &\leq \frac{t^2}{2}\|e_0\|^2 - \frac{t^4}{4} \int_{|x|\leq \frac{1}{4}} \int_{|y|\leq \frac{1}{4}} \ln \frac{1}{|x-y|} e_0^2(y)e_0^2(x)dydx \\
 &\leq \frac{t^2}{2}\|e_0\|^2 - \frac{t^4 \ln 2}{4} \left(\int_{\mathbb{R}^2} e_0^2(x)dx \right)^2,
 \end{aligned}
 \tag{3.7}$$

which implies that there exists $t_0 > 0$ large enough such that $I_{\alpha,\lambda}(t_0e_0) < 0$. □

Based on the mountain pass theorem without the Palais–Smale condition (see [43]), there exists a $(PS)_{c_{\alpha,\lambda}}$ sequence $\{u_n\} \subset H^1(\mathbb{R}^2)$, that is,

$$I_{\alpha,\lambda}(u_n) \rightarrow c_{\alpha,\lambda} \quad \text{and} \quad I'_{\alpha,\lambda}(u_n) \rightarrow 0.
 \tag{3.8}$$

Here $c_{\alpha,\lambda}$ is the mountain pass level characterized by

$$c_{\alpha,\lambda} = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} I_{\alpha,\lambda}(\gamma(t))
 \tag{3.9}$$

with

$$\Gamma := \{\gamma \in C^1([0, 1], H^1(\mathbb{R}^2)) : \gamma(0) = 0 \quad \text{and} \quad I_{\alpha,\lambda}(\gamma(1)) < 0\}.$$

Remark 3.2 Observe from Lemma 3.1 that there exist two constants $a, b > 0$ independently of α, λ such that $a < c_{\alpha,\lambda} < b$.

Lemma 3.3 *Let $\{u_n\} \subset H^1(\mathbb{R}^2)$ be a $(PS)_{c_{\alpha,\lambda}}$ sequence of $I_{\alpha,\lambda}$ for fixed $\alpha, \lambda \in (0, 1)$, then $\{u_n\}$ is bounded in $H^1(\mathbb{R}^2)$.*

Proof Observe that there exists $C_1, C_2 > 0$ such that

$$\begin{aligned}
 & C_1 + C_2 \|u_n\| \\
 & \geq I_{\alpha,\lambda}(u_n) - \frac{1}{4} I'_{\alpha,\lambda}(u_n) u_n \\
 & = \frac{1}{4} \|u_n\|^2 + \frac{3\lambda}{20} \|u_n\|_2^{\frac{5}{2}} + \int_{A_n \cup \{\mathbb{R}^2 \setminus A_n\}} \left(\frac{1}{4} f(u_n) u_n - F(u_n) \right) dx + \frac{r-4}{4r} \lambda \int_{\mathbb{R}^2} |u_n|^r dx \\
 & \geq \frac{1}{4} \|u_n\|^2 + \frac{3\lambda}{20} \|u_n\|_2^{\frac{5}{2}} - \int_{A_n} \left(F(u_n) - \frac{1}{4} f(u_n) u_n \right) dx + \frac{r-4}{4r} \lambda \int_{\mathbb{R}^2} |u_n|^r dx,
 \end{aligned} \tag{3.10}$$

where $A_n := \{x \mid \frac{1}{4} f(u_n) u_n - F(u_n) \leq 0\}$. Recalling (f_1) and (f_5) , we can take $t^* = \max\{t_0, 4M_0\}$ so that there exists $C_{t^*} > 0$ such that

$$\begin{aligned}
 & \int_{A_n} \left(F(u_n) - \frac{1}{4} f(u_n) u_n \right) dx \\
 & = \int_{A_n \cap \{|u_n| \leq 4t^*\}} \left(F(u_n) - \frac{1}{4} f(u_n) u_n \right) dx + \int_{A_n \cap \{|u_n| \geq 4t^*\}} \left(F(u_n) - \frac{1}{4} f(u_n) u_n \right) dx \\
 & \leq C_{t^*} \int_{A_n \cap \{|u_n| \leq 4t^*\}} u_n^2 dx + \int_{A_n \cap \{|u_n| \geq 4t^*\}} \left(M_0 - \frac{1}{4} u_n \right) f(u_n) dx \\
 & \leq C_{t^*} \|u_n\|_2^2.
 \end{aligned} \tag{3.11}$$

Since for any large $B_1 > 0$, there exists $B_2 > 0$ such that $\frac{3}{20} \|u_n\|_2^{\frac{5}{2}} \geq B_1 \|u_n\|_2^2 - B_2$. So combining (3.10) and (3.11) we have

$$C_1 + C_2 \|u_n\| + \lambda B_2 \geq \frac{1}{8} \|u_n\|^2 + \int_{\mathbb{R}^2} \left[(\lambda B_1 - C_{t^*}) |u_n|^2 + \frac{r-4}{4r} \lambda |u_n|^r \right] dx. \tag{3.12}$$

We can obtain

$$(\lambda B_1 - C_{t^*}) t^2 + \frac{r-4}{4r} \lambda t^r \geq 0$$

for $t \geq 0$, by letting B_1 can be chosen arbitrary large. Thus, it follows from (3.12) that $\|u_n\| \leq C$ for some C independently of n . \square

Let us define Moser type functions $w_n(x)$ supported in $B_\rho(0)$ as follows:

$$w_n(x) = \frac{1}{\sqrt{2\pi}} \begin{cases} \sqrt{\log n}, & 0 \leq |x| \leq \rho/n, \\ \frac{\log(\rho/|x|)}{\sqrt{\log n}}, & \rho/n \leq |x| \leq \rho, \\ 0, & |x| \geq \rho, \end{cases}$$

where ρ is given in (f_6) . An estimation yields

$$\begin{aligned}
 \|w_n\|^2 & = \int_{B_\rho(0)} (|\nabla w_n|^2 + w_n^2) dx \\
 & = 1 + \rho^2 \left(\frac{1}{4 \log n} - \frac{1}{4n^2 \log n} - \frac{1}{2n^2} \right) \\
 & =: 1 + \rho^2 \delta_n,
 \end{aligned} \tag{3.13}$$

and

$$\frac{2\lambda}{5} \|w_n\|_2^{\frac{5}{2}} = \frac{2\lambda}{5} \rho^{5/2} \left(\frac{1}{4 \log n} - \frac{1}{4n^2 \log n} - \frac{1}{2n^2} \right)^{5/4} = \frac{2\lambda}{5} \rho^{5/2} \delta_n^{5/4}. \tag{3.14}$$

Moreover,

$$\begin{aligned} \frac{1}{4} \int_{\mathbb{R}^2} (G_\alpha(x) * w_n^2) w_n^2 dx &= \frac{1}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|x-y|^{-\alpha} - 1}{\alpha} w_n^2(x) w_n^2(y) dx dy \\ &= \frac{1}{4} \int_{B_\rho(0)} \int_{B_\rho(0)} \frac{|x-y|^{-\alpha} - 1}{\alpha} w_n^2(x) w_n^2(y) dx dy \\ &\geq 0. \end{aligned} \tag{3.15}$$

Lemma 3.4 $\sup_{\alpha \in (0, 1), \lambda \in (0, \lambda^*)} c_{\alpha, \lambda} < \frac{2\pi}{\theta_0}$, where $\lambda^* := \min \left\{ 1, \sqrt[4]{\frac{\theta_0}{5\pi}} \right\}$.

Proof Recalling (f_6) , for

$$\varepsilon \in \left(0, \kappa - \frac{1}{4\rho^2 \sqrt{\pi \theta_0}} \right), \tag{3.16}$$

there exists $t_\varepsilon > 0$ such that,

$$t^2 F(t) \geq (\kappa - \varepsilon) e^{\theta_0 t^2}, \quad \text{for } t \geq t_\varepsilon. \tag{3.17}$$

We proceed the proof by considering three cases.

Case 1. $t \in \left[0, \sqrt{\frac{2\pi}{\theta_0}} \right]$, then by (3.13)-(3.15), we have for large n

$$\begin{aligned} &I_{\alpha, \lambda}(tw_n) \\ &\leq \frac{t^2}{2} \|w_n\|^2 - \frac{t^4}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|x-y|^{-\alpha} - 1}{\alpha} w_n^2(y) w_n^2(x) dy dx - \int_{\mathbb{R}^2} F(tw_n) dx + \frac{2}{5} \|tw_n\|_2^{\frac{5}{2}} \\ &\leq \frac{t^2}{2} \|w_n\|^2 + \frac{2t^{\frac{5}{2}}}{5} \|w_n\|_2^{\frac{5}{2}} \\ &\leq \frac{2\pi}{\theta_0} \left[\frac{1}{2} \|w_n\|^2 + \frac{2}{5} \sqrt{\frac{2\pi}{\theta_0}} \|w_n\|_2^{\frac{5}{2}} \right] \\ &< \frac{2\pi}{\theta_0}. \end{aligned} \tag{3.18}$$

Case 2. $t \in \left[\sqrt{\frac{5\pi}{\theta_0}}, +\infty \right)$. According to the definition of w_n , we have for large $n \in \mathbb{N}$, $tw_n(x) \geq t_\varepsilon$ for $x \in B_{\rho/n}$. From (3.13)–(3.17), we deduce that for large n ,

$$\begin{aligned}
 & I_{\alpha,\lambda}(tw_n) \\
 & \leq \frac{t^2}{2} \|w_n\|^2 - \frac{t^4}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|x-y|^{-\alpha} - 1}{\alpha} w_n^2(y) w_n^2(x) dy dx - \int_{\mathbb{R}^2} F(tw_n) dx + \frac{2}{5} \|tw_n\|_2^{\frac{5}{2}} \\
 & \leq \frac{t^2}{2} \|w_n\|^2 + \frac{2t^{\frac{5}{2}}}{5} \|w_n\|_2^{\frac{5}{2}} - \frac{2\pi^2(\kappa - \varepsilon)\rho^2}{n^2 t^2 \log n} e^{\frac{\theta_0 t^2 \log n}{2\pi}} \\
 & = \frac{1 + \delta_n \rho^2}{2} t^2 + \frac{2}{5} t^{\frac{5}{2}} \rho^{\frac{5}{2}} \delta_n^{\frac{5}{4}} - \frac{2\pi^2(\kappa - \varepsilon)\rho^2}{n^2 t^2 \log n} e^{\frac{\theta_0 t^2 \log n}{2\pi}} \\
 & < t^2 + \frac{2}{5} \rho^{\frac{5}{2}} t^{\frac{5}{2}} - \frac{2\pi^2(\kappa - \varepsilon)\rho^2}{n^2 t^2 \log n} e^{\frac{\theta_0 t^2 \log n}{2\pi}}.
 \end{aligned} \tag{3.19}$$

Define

$$g(n) := \frac{2\pi^2(\kappa - \varepsilon)\rho^2}{n^2 t^2 \log n} e^{\frac{\theta_0 t^2 \log n}{2\pi}} = \frac{2\pi^2(\kappa - \varepsilon)\rho^2}{n^2 t^2 \log n} n^{\frac{\theta_0}{2\pi} t^2},$$

then

$$g(n) \geq \frac{2\pi^2(\kappa - \varepsilon)\rho^2}{4t^2 \log 2} 2^{\frac{\theta_0}{2\pi} t^2} \tag{3.20}$$

for large n . It then follows from (3.19) and (3.20) that

$$\begin{aligned}
 I_{\alpha,\lambda}(tw_n) & < t^2 + \frac{2}{5} \rho^{\frac{5}{2}} t^{\frac{5}{2}} - \frac{2\pi^2(\kappa - \varepsilon)\rho^2}{4t^2 \log 2} 2^{\frac{\theta_0}{2\pi} t^2} \\
 & \rightarrow -\infty, \text{ as } t \rightarrow \infty,
 \end{aligned} \tag{3.21}$$

which implies that

$$I_{\alpha,\lambda}(tw_n) < \frac{2\pi}{\theta_0}. \tag{3.22}$$

Case 3. $t \in \left[\sqrt{\frac{2\pi}{\theta_0}}, \sqrt{\frac{5\pi}{\theta_0}} \right]$. According to the definition of w_n , we have for large $n \in \mathbb{N}$, $tw_n(x) \geq t_\varepsilon$ for $x \in B_{\rho/n}$. Since $\lambda \in \left(0, \min \left\{ 1, \sqrt{\frac{4\theta_0}{5\pi}} \right\} \right)$, it then follows from (3.13)–(3.17) that for large n

$$\begin{aligned}
 & I_{\alpha,\lambda}(tw_n) \\
 & \leq \frac{t^2}{2} \|w_n\|^2 - \frac{t^4}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|x-y|^{-\alpha} - 1}{\alpha} w_n^2(y) w_n^2(x) dy dx - \int_{\mathbb{R}^2} F(tw_n) dx + \frac{2\lambda}{5} \|tw_n\|_2^{\frac{5}{2}} \\
 & < \frac{t^2}{2} \|w_n\|^2 + \frac{2\lambda t^{\frac{5}{2}}}{5} \|w_n\|_2^{\frac{5}{2}} - \frac{2\pi^2(\kappa - \varepsilon)\rho^2}{n^2 t^2 \log n} e^{\frac{\theta_0 t^2 \log n}{2\pi}} \\
 & < \frac{t^2}{2} \|w_n\|^2 + \frac{2\lambda t^{\frac{5}{2}}}{5} \|w_n\|_2^{\frac{5}{2}} - \frac{2\pi\theta_0(\kappa - \varepsilon)\rho^2}{5n^2 \log n} e^{\frac{\theta_0 t^2 \log n}{2\pi}} \\
 & < \frac{t^2}{2} \|w_n\|^2 + \frac{2t^2}{5} \|w_n\|_2^{\frac{5}{2}} - \frac{2\pi\theta_0(\kappa - \varepsilon)\rho^2}{5n^2 \log n} e^{\frac{\theta_0 t^2 \log n}{2\pi}} := \psi_n(t).
 \end{aligned} \tag{3.23}$$

Then there exists $t_n > 0$ such that $\psi_n(t_n) = \max_{t>0} \psi_n(t)$ and

$$t_n^2 = \frac{4\pi}{\theta_0} \left[1 + \frac{\log \left(1 + \rho^2 \delta_n + \frac{4}{5} \rho^{\frac{5}{2}} \delta_n^{\frac{5}{4}} \right) - \log (2\pi \theta_0 (\kappa - \varepsilon) \rho^2) + 2 \log \sqrt{\frac{5\pi}{\theta_0}}}{2 \log n} \right]. \tag{3.24}$$

Obviously, we can easily see that $t_n \in \left[\sqrt{\frac{2\pi}{\theta_0}}, \sqrt{\frac{5\pi}{\theta_0}} \right]$ for large n . Then, by (3.23) and (3.24) we have

$$\begin{aligned} \psi_n(t) &\leq \psi_n(t_n) \\ &= \left(\frac{1}{2} \|w_n\|^2 + \frac{2}{5} \|w_n\|^{\frac{5}{2}} \right) t_n^2 - \frac{\pi \left(\|w_n\|^2 + \frac{4}{5} \|w_n\|^{\frac{5}{2}} \right)}{\theta_0 \log n} \\ &= \frac{4\pi}{\theta_0} \left(\frac{1}{2} + \frac{\rho^2 \delta_n}{2} + \frac{2}{5} \rho^{\frac{5}{2}} \delta_n^{\frac{5}{4}} \right) \left[1 + \frac{\log \left(1 + \rho^2 \delta_n + \frac{4}{5} \rho^{\frac{5}{2}} \delta_n^{\frac{5}{4}} \right)}{2 \log n} \right. \\ &\quad \left. - \frac{\log \left(2\theta_0 (\kappa - \varepsilon) \rho^2 \sqrt{\frac{5\pi}{\theta_0}} \right)}{2 \log n} - \frac{1}{2 \log n} \right] \\ &\leq \frac{4\pi}{\theta_0} \left(\frac{1}{2} + \frac{\rho^2 \delta_n}{2} + \frac{2}{5} \rho^{\frac{5}{2}} \delta_n^{\frac{5}{4}} \right) \left[1 + \frac{\log \left(1 + \rho^2 \delta_n + \frac{4}{5} \rho^{\frac{5}{2}} \delta_n^{\frac{5}{4}} \right)}{2 \log n} - \frac{1}{2 \log n} \right], \end{aligned} \tag{3.25}$$

which, together with the definition of δ_n , implies that

$$\begin{aligned} \psi_n(t) &\leq \frac{4\pi}{\theta_0} \left[\frac{1}{2} + \frac{\rho^2}{8 \log n} + \frac{o(1)}{\log n} - \frac{1}{4 \log n} \right] \\ &< \frac{2\pi}{\theta_0}, \text{ for } n \text{ large.} \end{aligned} \tag{3.26}$$

Combining (3.18), (3.22) and (3.26), we have $I_{\alpha,\lambda}(t w_n) < \frac{2\pi}{\theta_0}$. It follows from (3.21) that, for fixed n large enough, there exists $t_0 > 0$ such that $I_{\alpha,\lambda}(t_0 w_n) < 0$. Define $\gamma(t) = t t_0 w_n$ for $t \in [0, 1]$, then $\gamma \in \Gamma$. Therefore, the conclusion follows immediately. The proof is complete. \square

In the following, we establish a critical exponential version of splitting lemma to the Palais–Smale sequences of $I_{\alpha,\lambda}$.

Lemma 3.5 *Assume that $\{u_n\}$ is a bounded $(PS)_{c_{\alpha,\lambda}}$ sequence of $I_{\alpha,\lambda}$ for fixed α and λ . Then there exist $B \in \mathbb{R}$ and a number $k \in \mathbb{N} \cup \{0\}$, and a finite sequence*

$$(u_0, w^1, \dots, w^k) \subset H^1(\mathbb{R}^2), \quad w^j > 0, \quad \text{for } j = 1, \dots, k \text{ (if } k \geq 1)$$

of critical points for the following functional

$$J_{B,\alpha,\lambda}(u) := \frac{1}{2}\|u\|^2 + \frac{B^2}{2\alpha} \int_{\mathbb{R}^2} u^2 dx + \frac{\lambda B^{1/2}}{2} \int_{\mathbb{R}^2} u^2 dx - \int_{\mathbb{R}^2} F(u) dx - \frac{1}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u^2 \right) u^2 dx - \frac{\lambda}{r} \|u\|_r^r.$$

and k sequences of points $\{y_n^j\} \subset \mathbb{R}^2$, $1 \leq j \leq k$, such that

- (i) $|y_n^j| \rightarrow +\infty$, $|y_n^j - y_n^i| \rightarrow +\infty$ if $i \neq j$, $n \rightarrow +\infty$,
- (ii) $\|u_n\|^2 = \|u_0\|^2 + \sum_{j=1}^k \|w^j\|^2 + o(1)$,
- (iii) $c_{\alpha,\lambda} + \frac{B^4}{4\alpha} + \frac{B^{5/2}\lambda}{10} = J_{B,\alpha,\lambda}(u_0) + \sum_{j=1}^k J_{B,\alpha,\lambda}(w^j)$,
- (iv) $B^2 = \|u_0\|_2^2 + \sum_{j=1}^k \|w^j\|_2^2$.

Otherwise, if $k = 0$, then $u_n \rightarrow u_0$ in $H^1(\mathbb{R}^2)$.

Proof Recall that $\{u_n\}$ is a bounded sequence in $H^1(\mathbb{R}^2)$, then, up to subsequence, there exist $u_0 \in H^1(\mathbb{R}^2)$ and $B \in \mathbb{R}$ such that $u_n \rightharpoonup u_0$ weakly in $H^1(\mathbb{R}^2)$ and $\int_{\mathbb{R}^2} |u_n|^2 dx \rightarrow B^2$ as $n \rightarrow \infty$. According to Remark 3.2, it follows from (3.10) and (3.11) that there exists $C_1 > 0$ such that

$$\int_{\tilde{A}_n} \left(\frac{1}{4} f(u_n)u_n - F(u_n) \right) dx \leq C_1, \tag{3.27}$$

where $\tilde{A}_n := \{x \mid \frac{1}{4} f(u_n)u_n - F(u_n) \geq 0\}$. Let $M > \max\{20M_0, t_0\}$, then from (f_5) and (3.27), we conclude that

$$\begin{aligned} & \frac{1}{5} \int_{|u_n| \geq M} f(u_n)u_n dx \\ & \leq \int_{|u_n| \geq M} \frac{1}{4} f(u_n)u_n - M_0 |f(u_n)| dx \\ & \leq \int_{|u_n| \geq M} \frac{1}{4} f(u_n)u_n - F(u_n) dx \\ & \leq \int_{\tilde{A}_n} \frac{1}{4} f(u_n)u_n - F(u_n) dx \\ & < C_1, \end{aligned} \tag{3.28}$$

which implies by (f_1) that $f(u_n)u_n$ is bounded in $L^1(\mathbb{R}^2)$. So we can infer that $f(u_n) \rightarrow f(u_0)$ in $L^1(B_R(0))$ for any $R > 0$ and then (see [24])

$$\int_{\mathbb{R}^2} f(u_n)\varphi dx \rightarrow \int_{\mathbb{R}^2} f(u_0)\varphi dx, \quad \varphi \in C_0^\infty(\mathbb{R}^2). \tag{3.29}$$

Then $J'_{B,\alpha,\lambda}(u_0) = 0$ for any fixed α, λ . Observe that

$$I_{\alpha,\lambda}(u_n) = J_{B,\alpha,\lambda}(u_n) - \frac{B^4}{4\alpha} - \frac{B^{5/2}\lambda}{10} + o(1),$$

and so $J_{B,\alpha,\lambda}(u_n) \rightarrow c_{\alpha,\lambda} + \frac{B^4}{4\alpha} + \frac{B^{5/2}\lambda}{10}$ and $J'_{B,\alpha,\lambda}(u_n) \rightarrow 0$ in H^{-1} as $n \rightarrow \infty$. Moreover, $v_n^1 \rightarrow 0$ in $H^1(\mathbb{R}^2)$ if we define $v_n^1 := u_n - u_0$. It follows from the Brezis-Lieb lemma [10] that

$$\begin{aligned} \|u_n\|^2 &= \|v_n^1\|^2 + \|u_0\|^2 + o(1), \\ \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_n^2 \right) u_n^2 dx &= \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_0^2 \right) u_0^2 dx + \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (v_n^1)^2 \right) (v_n^1)^2 dx + o(1), \end{aligned} \tag{3.30}$$

where the second identity can be proved similarly as that of Lemma 2.2 in [25] and we omit it. Using (f₀) and (f₅), we can also verify

$$\int_{\mathbb{R}^2} F(u_n) dx = \int_{\mathbb{R}^2} F(u_0) dx + \int_{\mathbb{R}^2} F(v_n^1) dx + o(1). \tag{3.31}$$

Combining (3.31) with (3.30), we immediately get

$$\begin{aligned} &J_{B,\alpha,\lambda}(u_n) - J_{B,\alpha,\lambda}(u_0) \\ &= \frac{1}{2} \|v_n^1\|^2 + \frac{B^2}{2\alpha} \int_{\mathbb{R}^2} |v_n^1|^2 dx + \frac{\lambda B^{1/2}}{2} \int_{\mathbb{R}^2} |v_n^1|^2 dx, \\ &\quad - \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (v_n^1)^2 \right) (v_n^1)^2 dx - \frac{\lambda}{r} \|v_n^1\|_r^r - \int_{\mathbb{R}^2} F(v_n^1) dx + o(1) \\ &= J_{B,\alpha,\lambda}(v_n^1) + o(1). \end{aligned} \tag{3.32}$$

Recalling $J'_{B,\alpha,\lambda}(u_0) = 0$ whose corresponding Pohozaev identity is

$$\begin{aligned} P_{B,\alpha,\lambda}(u_0) &:= \left(1 + \frac{B^2}{\alpha} + \lambda B^{1/2} \right) \|u_0\|_2^2 - \frac{4-\alpha}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (u_0)^2 \right) (u_0)^2 dx \\ &\quad - 2 \int_{\mathbb{R}^2} \left(F(u_0) + \frac{\lambda}{r} |u_0|^r \right) dx = 0, \end{aligned}$$

we can define

$$\begin{aligned} \mathcal{B}_{B,\alpha,\lambda}(u_0) &:= 2J'_{B,\alpha,\lambda}(u_0)u_0 - P_{B,\alpha,\lambda}(u_0) \\ &= 2\|\nabla u_0\|_2^2 + \left(1 + \frac{B^2}{\alpha} + \lambda B^{1/2} \right) \|u_0\|_2^2 - \frac{4+\alpha}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_0^2 \right) u_0^2 dx \\ &\quad - \left(2 - \frac{2}{r} \right) \lambda \|u_0\|_r^r + 2 \int_{\mathbb{R}^2} [F(u_0) - f(u_0)u_0] dx = 0, \end{aligned}$$

which yields that

$$\begin{aligned} 4J_{B,\alpha,\lambda}(u_0) &= 4J_{B,\alpha,\lambda}(u_0) - \mathcal{B}_{B,\alpha,\lambda}(u_0) \\ &= \left(1 + \frac{B^2}{\alpha} + \lambda B^{1/2} \right) \|u_0\|_2^2 + \frac{1}{4} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_0^2 \right) u_0^2 dx \\ &\quad + 2 \int_{\mathbb{R}^2} [f(u_0)u_0 - 3F(u_0)] dx + \left(2 - \frac{6}{r} \right) \lambda \|u_0\|_r^r. \end{aligned} \tag{3.33}$$

Consider sequence $\{v_n^1\}$. We claim that either

- (v1) $v_n^1 \rightarrow 0$ in $H^1(\mathbb{R}^2)$ as $n \rightarrow \infty$, or

(v2) there exist $m > 0$ and $\{y_n^1\} \subset \mathbb{R}^2$ such that

$$\liminf_{n \rightarrow \infty} \int_{B_1(y_n^1)} |v_n^1|^2 dx \geq m > 0. \tag{3.34}$$

If (3.34) does not occur, then by Lions’ vanishing lemma (see [30, 31]), we have $v_n^1 \rightarrow 0$ in $L^s(\mathbb{R}^2)$ for all $s > 2$. And so, using the Hardy–Littlewood–Sobolev inequality, we have $\int_{\mathbb{R}^2} (\frac{1}{|x|^\alpha} * (v_n^1)^2)(v_n^1)^2 dx = o(1)$. Moreover, arguing similarly as in [24], we have $\int_{\mathbb{R}^2} F(v_n^1) dx = o(1)$. Here we consider two cases:

Case 1. $u_0 \equiv 0$. That is, $v_n^1 \equiv u_n$. Combining (3.32) with (3.33) we have

$$\begin{aligned} J_{B,\alpha,\lambda}(u_n) &= \frac{1}{2} \|u_n\|^2 + \frac{B^2}{2\alpha} \int_{\mathbb{R}^2} |u_n|^2 dx + \frac{\lambda B^{1/2}}{2} \int_{\mathbb{R}^2} |u_n|^2 dx + o(1) \\ &\geq \frac{1}{2} \|u_n\|^2 + \frac{B^4}{2\alpha} + \frac{\lambda B^{5/2}}{2} + o(1), \end{aligned} \tag{3.35}$$

which implies by $c_{\alpha,\lambda} < \frac{2\pi}{\theta_0}$ that $\sup_{n \in \mathbb{N}} \|u_n\|^2 < \frac{4\pi}{\theta_0}$. Then there exists $\varepsilon_0 > 0$ small such that

$$\|u_n\|^2 < \frac{4\pi}{\theta_0} (1 - 5\varepsilon_0), \tag{3.36}$$

and then there exists $s \in (1, 2)$ such that

$$(1 + \varepsilon_0)(1 - 5\varepsilon_0)s < 1.$$

For any $\xi > 0$, there exists $C_\xi > 0$ such that

$$|f(t)|^s \leq \xi |t| + C_\xi [e^{\theta_0(1+\varepsilon_0)t^2} - 1], \quad t \geq 0. \tag{3.37}$$

From Lemma 1.1, (3.36) and (3.37), Hölder’s inequality, we deduce that

$$\begin{aligned} \int_{\mathbb{R}^2} f(u_n)u_n dx &\leq \xi \|u_n\|^2 + C_\xi \left(\int_{\mathbb{R}^2} |f(u_n)|^s dx \right)^{1/s} \|u_n\|_{s'} \\ &\leq \xi \|u_n\|^2 + C_\xi \left(\int_{\mathbb{R}^2} [e^{\theta_0(1+\varepsilon_0)s|u_n|^2} - 1] dx \right)^{1/s} \|u_n\|_{s'} \\ &\leq \xi \|u_n\|^2 + C_\xi \left(\int_{\mathbb{R}^2} [e^{\|u_n\|^2 \theta_0(1+\varepsilon_0)s \frac{|u_n|^2}{\|u_n\|^2}} - 1] dx \right)^{1/s} \|u_n\|_{s'} \\ &\leq \xi \|u_n\|^2 + C_\xi \|u_n\|_{s'} = o(1). \end{aligned} \tag{3.38}$$

Here, $s' = \frac{s}{s-1} \in (2, +\infty)$. Thus,

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^2} f(u_n)u_n dx = 0. \tag{3.39}$$

From $J'_{\alpha,B,\lambda}(u_n)u_n = o(1)$, we deduce that $u_n \rightarrow 0$ in $H^1(\mathbb{R}^2)$ which contradicts the fact that

$$J_{B,\alpha,\lambda}(u_n) \rightarrow c_{\alpha,\lambda} + \frac{B^4}{4\alpha} + \frac{B^{5/2}\lambda}{10}$$

as $n \rightarrow \infty$. Thus (v2) holds true for $\{v_n^1\}$.

Case 2. $u_0 \neq 0$. That is, $\|u_0\| > 0$. In order to prove that (v1) holds for $\{v_n^1\}$, and Lemma 3.5 holds with $k = 0$, we only need to show that

$$\|u_n\|^2 \rightarrow \|u_0\|^2 \text{ as } n \rightarrow \infty. \tag{3.40}$$

Indeed, by Fatou’s lemma, we have

$$\begin{aligned} I_{\alpha,\lambda}(u_0) &= \frac{1}{2}\|u_0\|^2 + \frac{1}{4\alpha}\|u_0\|_2^2 + \frac{2\lambda}{5}\|u_0\|_2^4 \\ &\quad - \frac{1}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (u_0)^2 \right) (u_0)^2 dx - \int_{\mathbb{R}^2} (F(u_0) + \frac{\lambda}{r}|u_0|^r) dx \\ &\leq \liminf_{n \rightarrow \infty} \left(\frac{1}{2}\|u_n\|^2 + \frac{1}{4\alpha}\|u_n\|_2^2 + \frac{2\lambda}{5}\|u_n\|_2^4 \right. \\ &\quad \left. - \frac{1}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (u_n)^2 \right) (u_n)^2 dx - \int_{\mathbb{R}^2} \left(F(u_n) + \frac{\lambda}{r}|u_n|^r \right) dx \right) \\ &= \liminf_{n \rightarrow \infty} I_{\alpha,\lambda}(u_n) = c_{\alpha,\lambda}. \end{aligned} \tag{3.41}$$

If $I_{\alpha,\lambda}(u_0) = c_{\alpha,\lambda}$, by (3.41) we obtain immediately (3.40) holds true. Otherwise if $I_{\alpha,\lambda}(u_0) < c_{\alpha,\lambda}$, then we have

$$\begin{aligned} \|u_0\|^2 + \frac{1}{2\alpha}\|u_0\|_2^4 + \frac{4\lambda}{5}\|u_0\|_2^5 \\ < 2c_{\alpha,\lambda} + 2 \int_{\mathbb{R}^2} F(u_0) dx + \frac{1}{2\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_0^2 \right) u_0^2 dx + \frac{2\lambda}{r}\|u_0\|_r^r. \end{aligned} \tag{3.42}$$

In view of the definition of $I_{\alpha,\lambda}$, we also have

$$\begin{aligned} \lim_{n \rightarrow \infty} \left(\|u_n\|^2 + \frac{1}{2\alpha}\|u_n\|_2^4 + \frac{4\lambda}{5}\|u_n\|_2^5 \right) \\ = 2c_{\alpha,\lambda} + 2 \int_{\mathbb{R}^2} F(u_0) dx + \frac{1}{2\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_0^2 \right) u_0^2 dx + \frac{2\lambda}{r}\|u_0\|_r^r. \end{aligned} \tag{3.43}$$

Take

$$w_n = \frac{u_n}{\left(\|u_n\|^2 + \frac{1}{2\alpha}\|u_n\|_2^4 + \frac{4\lambda}{5}\|u_n\|_2^5 \right)^{1/2}}$$

and

$$w_0 = \frac{u_0}{\sqrt{2c_{\alpha,\lambda} + 2 \int_{\mathbb{R}^2} F(u_0) dx + \frac{1}{2\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_0^2 \right) u_0^2 dx + \frac{2\lambda}{r}\|u_0\|_r^r}}.$$

It then follows from (3.42) and (3.43) that $\|w_n\| \leq 1$, $w_n \rightharpoonup w_0$, and $\|w_0\| < 1$. Similarly to Lions [32], one has that

$$\sup_{n \in \mathbb{N}} \left(\int_{\mathbb{R}^2} e^{4\pi p w_n^2} - 1 \right) dx < \infty \tag{3.44}$$

for all

$$p < \bar{p} := \frac{1}{A - \|w_0\|^2} = 2 \frac{c_{\alpha,\lambda} + \int_{\mathbb{R}^2} F(u_0) dx + \frac{1}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_0^2 \right) u_0^2 dx + \frac{\lambda}{r}\|u_0\|_r^r}{\|u_n\|^2 - \|u_0\|^2} + o(1),$$

where $A = \lim_{n \rightarrow \infty} \|w_n\|^2$. Recalling (3.32) with (3.33), we have

$$\begin{aligned}
 c_{\alpha,\lambda} + \frac{B^4}{4\alpha} + \frac{B^{5/2}\lambda}{10} &= J_{B,\alpha,\lambda}(u_n) + o(1) \\
 &= J_{B,\alpha,\lambda}(u_0) + \frac{1}{2}\|v_n^1\|^2 + \frac{B^2}{2\alpha} \int_{\mathbb{R}^2} |v_n^1|^2 dx + \frac{\lambda B^{1/2}}{2} \int_{\mathbb{R}^2} |v_n^1|^2 dx + o(1) \\
 &\geq \frac{1}{4} \left(\frac{B^2}{\alpha} + \lambda B^{1/2} \right) \|u_0\|_2^2 + \frac{1}{2}\|v_n^1\|^2 + \frac{B^2}{2\alpha} \int_{\mathbb{R}^2} |v_n^1|^2 dx + \frac{\lambda B^{1/2}}{2} \int_{\mathbb{R}^2} |v_n^1|^2 dx + o(1) \\
 &\geq \frac{1}{4} \left(\frac{B^2}{\alpha} + \lambda B^{1/2} \right) \|u_n\|_2^2 + \frac{1}{2}\|v_n^1\|^2 + o(1), \\
 &= \frac{1}{4} \left(\frac{B^4}{\alpha} + \lambda B^{5/2} \right) + \frac{1}{2}\|v_n^1\|^2 + o(1),
 \end{aligned}
 \tag{3.45}$$

which implies by $c_{\alpha,\lambda} < \frac{2\pi}{\theta_0}$ that $\sup_{n \in \mathbb{N}} \|v_n^1\|^2 < \frac{4\pi}{\theta_0}$. And so,

$$\frac{\theta_0}{2\pi} < \frac{2}{D - \|u_0\|^2} = \lim_{n \rightarrow \infty} \frac{2}{\|v_n^1\|^2},$$

where $D = \lim_{n \rightarrow \infty} \|u_n\|^2$. Then recalling (3.43), we can always choose $q > 1$ sufficiently close to 1 and $\varepsilon > 0$ small such that

$$\begin{aligned}
 q(\theta_0 + \varepsilon) \left(\|u_n\|^2 + \frac{1}{2\alpha} \|u_n\|_2^4 + \frac{4\lambda}{5} \|u_n\|_2^5 \right) \\
 \leq 4\pi p < 4\pi \frac{1}{A - \|u_0\|^2} \\
 = 8\pi \frac{c_{\alpha,\lambda} + \int_{\mathbb{R}^2} F(u_0) dx + \frac{1}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_0^2 \right) u_0^2 dx + \frac{\lambda}{r} \|u_0\|^r}{\|u_n\|^2 - \|u_0\|^2} + o(1)
 \end{aligned}$$

for some p satisfying (3.44). Based on the above facts, using condition (f_0) , we have

$$\int_{\mathbb{R}^2} |f(u_n)|^q dx \leq C \|u_n\|_q^q + C \int_{\mathbb{R}^2} \left[e^{q(\theta_0 + \varepsilon)} \left(\|u_n\|^2 + \frac{1}{2\alpha} \|u_n\|_2^4 + \frac{4\lambda}{5} \|u_n\|_2^5 \right) w_n^2 - 1 \right] dx < C$$

for some $C > 0$ independently of n . In virtue of the above facts and (3.29), we have

$$\begin{aligned}
 &\left| \int_{\mathbb{R}^2} f(u_n)u_n - f(u_0)u_0 dx \right| \\
 &= \left| \int_{\mathbb{R}^2} f(u_n)(u_n - u_0) - (f(u_n) - f(u_0))u_0 dx \right| \\
 &\leq \left(\int_{\mathbb{R}^2} |f(u_n)|^q dx \right)^{\frac{1}{q}} \left(\int_{\mathbb{R}^2} |u_n - u_0|^{\frac{q}{q-1}} dx \right)^{\frac{q-1}{q}} + \left| \int_{\mathbb{R}^2} (f(u_n) - f(u_0))u_0 dx \right| \\
 &\rightarrow 0, \quad \text{as } n \rightarrow \infty.
 \end{aligned}$$

By the fact that $(J'_{B,\alpha,\lambda}(u_n) - J'_{B,\alpha,\lambda}(u_0))(u_n - u_0) = o(1)$, we have $\|u_n - u_0\| = o(1)$, which implies that $I_{\alpha,\lambda}(u_0) = c_{\alpha,\lambda}$. This is a contradiction. Hence, $\|v_n^1\| \rightarrow 0$ and (v1) holds for $\{v_n^1\}$, Lemma 3.5 holds with $k = 0$.

If (v2) holds, namely, (3.34) is true, then there exists $w^1 \in H^1(\mathbb{R}^2) \setminus \{0\}$ such that $v_n^1(\cdot + y_n^1) \rightharpoonup w^1$ in $H^1(\mathbb{R}^2)$ and $u_n(\cdot + y_n^1) \rightharpoonup w^1$ in $H^1(\mathbb{R}^2)$. Recalling the fact that $v_n^1 \rightarrow 0$ in $H^1(\mathbb{R}^2)$, we find that $\{y_n^1\}$ must be unbounded. That is, $|y_n^1| \rightarrow +\infty$. Let us now show

that $J'_{B,\alpha,\lambda}(w^1) = 0$. Indeed, it suffices to show that $J'_{B,\alpha,\lambda}(u_n(\cdot + y_n^1))\varphi \rightarrow 0$ for fixed $\varphi \in C^\infty_0(\mathbb{R}^2)$. Since $J'_{B,\alpha,\lambda}(u_n) \rightarrow 0$ in H^{-1} as $n \rightarrow \infty$, and then $J'_{B,\alpha,\lambda}(u_n)\varphi(\cdot - y_n^1) \rightarrow 0$ for any $\varphi \in C^\infty_0(\mathbb{R}^2)$. Thus, it follows that as $n \rightarrow \infty$,

$$\begin{aligned} & J'_{B,\alpha,\lambda}(u_n(\cdot + y_n^1))\varphi \\ &= \int_{\mathbb{R}^2} [\nabla u_n(x + y_n^1)\nabla\varphi + u_n(x + y_n^1)\varphi]dx + \left(\frac{B^2}{\alpha} + \lambda B^{1/2}\right) \int_{\mathbb{R}^2} u_n(x + y_n^1)\varphi dx \\ &\quad - \frac{1}{\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x + y_n^1|^\alpha} * u_n^2\right) u_n(x + y_n^1)\varphi dx - \lambda \int_{\mathbb{R}^2} |u_n(\cdot + y_n^1)|^{r-2} u_n(\cdot + y_n^1)\varphi dx \\ &\quad - \int_{\mathbb{R}^2} f(u_n(\cdot + y_n^1))\varphi dx \rightarrow 0. \end{aligned}$$

So, $J'_{B,\alpha,\lambda}(w^1) = 0$. Set

$$v_n^2(x) = v_n^1(x) - w^1(x - y_n^1), \tag{3.46}$$

then using the fact that $v_n^1(\cdot + y_n^1) \rightarrow w^1$ in $H^1(\mathbb{R}^2)$, we have $v_n^2 \rightarrow 0$ in $H^1(\mathbb{R}^2)$. Using again the Brezis-Lieb lemma that

$$\begin{aligned} \|u_n\|^2 &= \|w^1\|^2 + \|u_0\|^2 + \|v_n^2\|^2 + o(1), \\ \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_n^2\right) u_n^2 dx &= \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u_0^2\right) u_0^2 dx + \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (w^1)^2\right) (w^1)^2 dx \\ &\quad + \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (v_n^2)^2\right) (v_n^2)^2 dx + o(1), \\ \int_{\mathbb{R}^2} F(u_n) dx &= \int_{\mathbb{R}^2} F(u_0) dx + \int_{\mathbb{R}^2} F(w^1) dx + \int_{\mathbb{R}^2} F(v_n^2) dx + o(1). \end{aligned} \tag{3.47}$$

By virtue of the above estimates, we deduce that

$$J_{B,\alpha,\lambda}(v_n^2) = J_{B,\alpha,\lambda}(u_n) - J_{B,\alpha,\lambda}(u_0) - J_{B,\alpha,\lambda}(w^1) + o(1). \tag{3.48}$$

Let us now study $\{v_n^2\}$. Since $\{v_n^2\}$ is bounded in $H^1(\mathbb{R}^2)$, one of (v1) and (v2) holds for $\{v_n^2\}$. The similar arguments used before imply that Lemma 3.5 holds with $k = 1$ if $v_n^2 \rightarrow 0$ in $H^1(\mathbb{R}^2)$. Otherwise, (v2) holds for $\{v_n^2\}$. We repeat the arguments above. Iterating this procedure, there exists sequence $\{y_n^j\} \subset \mathbb{R}^2$ such that $|y_n^j| \rightarrow +\infty$, $|y_n^j - y_n^i| \rightarrow +\infty$ if $i \neq j$ as $n \rightarrow +\infty$ and $v_n^j = v_n^{j-1} - w^{j-1}(x - y_n^{j-1})$ (like (3.46)) with $j \geq 2$ such that

$$v_n^j \rightarrow 0 \text{ in } H^1(\mathbb{R}^2), \quad J'_{B,\alpha,\lambda}(w^j) = 0.$$

Moreover, by the properties of the weak convergence, we get

$$\begin{aligned} (a) \quad & \|u_n\|^2 - \|u_0\|^2 - \sum_{i=1}^{j-1} \|w^i\|^2 = \|u_n - u_0 - \sum_{i=1}^j w^i(\cdot - y_n^i)\|^2 + o(1), \\ (b) \quad & c_{\alpha,\lambda} + \frac{B^4}{4\alpha} = J_{\alpha,B,\lambda}(u_0) + \sum_{i=1}^{j-1} J_{\alpha,B,\lambda}(w^i) + J_{\alpha,B,\lambda}(v_n^j) + o(1), \\ (c) \quad & B^2 = \|u_0\|_2^2 + \sum_{i=1}^{j-1} \|w^i\|_2^2 + \|v_n^{j+1}\|_2^2 + o(1). \end{aligned} \tag{3.49}$$

Now, we claim that there exists $C > 0$ such that $\|w^i\|^2 \geq C$, $i = 1, 2, \dots, k$. Without loss of generality, we can assume that $\|w^i\|^2 < \frac{2\pi}{\theta_0}$ for some i . Using (3.4), (3.5), $J'_{B,\alpha,\lambda}(w^i)w^i = 0$, the Hardy–Littlewood–Sobolev inequality, the Moser–Trudinger inequality and Lemma 2.3, one finds that for any $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that

$$\begin{aligned} \|w^i\|^2 &\leq \int \int_{|x-y|\leq 1} \frac{|x-y|^{-\alpha}-1}{\alpha} |w^i(x)|^2 |w^i(y)|^2 dx dy + C \|w^i\|^3 + C \|w^i\|^r \\ &\leq \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|w^i(x)|^2 |w^i(y)|^2}{|x-y|} dx dy + C \|w^i\|^3 + C \|w^i\|^r \\ &\leq C \|w^i\|^4 + C \|w^i\|^3 + C \|w^i\|^r. \end{aligned}$$

Hence, the claim is true. Recall that $\{u_n\}$ is bounded in $H^1(\mathbb{R}^2)$, from (3.49)(a) we deduce that the iteration must stop at some finite index k . And so $v_n^{k+1} \rightarrow 0$ in $H^1(\mathbb{R}^2)$ as $n \rightarrow \infty$. The proof is complete. \square

If $u \in H^1(\mathbb{R}^2)$ is a critical point of $I_{\alpha,\lambda}$, we have

$$\begin{aligned} \mathcal{B}_{\alpha,\lambda}(u) &:= 2\|\nabla u\|_2^2 + \|u\|_2^2 + \frac{1}{\alpha} \|u\|_2^4 + \lambda \|u\|_2^{5/2} - \frac{4+\alpha}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u^2\right) u^2 dx \\ &\quad + 2 \int_{\mathbb{R}^2} [F(u) - f(u)u] dx - \left(2 - \frac{2}{r}\right) \lambda \|u\|_r^r = 0, \end{aligned}$$

since $\mathcal{B}_{\alpha,\lambda}(u) = 2I'_{\alpha,\lambda}(u)u - P_{\alpha,\lambda}(u)$. Here, $P_{\alpha,\lambda}(u)$ is the associated Pohozaev functional with $I'_{\alpha,\lambda}(u) = 0$.

Lemma 3.6 *If $u \in H^1(\mathbb{R}^2) \setminus \{0\}$ satisfies $\mathcal{B}_{\alpha,\lambda}(u) = 0$, then there exists $\gamma \in C([0, 1], H^1(\mathbb{R}^2))$ such that $\gamma(0) = 0$, $I_{\alpha,\lambda}(\gamma(1)) < 0$, $u \in \gamma([0, 1])$, $0 \notin \gamma((0, 1])$ and*

$$\max_{t \in [0, 1]} I_{\alpha,\lambda}(\gamma(t)) = I_{\alpha,\lambda}(u).$$

Proof For $t \in (0, \infty)$, define $u_t := t^2 u(t \cdot)$, then we have

$$\begin{aligned} g(t) := I_{\alpha,\lambda}(u_t) &= \frac{t^4}{2} \|\nabla u\|_2^2 + \frac{t^2}{2} \|u\|_2^2 + \frac{t^4}{4\alpha} \|u\|_2^4 - \frac{t^{4+\alpha}}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u^2\right) u^2 dx \\ &\quad + \frac{2\lambda}{5} t^{5/2} \|u\|_2^{5/2} - \frac{\lambda t^{2(r-1)}}{r} \|u\|_r^r - t^{-2} \int_{\mathbb{R}^2} F(t^2 u) dx, \end{aligned}$$

which implies that for $t > 0$ large enough

$$I_{\alpha,\lambda}(u_t) < 0.$$

That is, there exists $t_0 > 1$ such that $I_{\alpha,\lambda}(u_{t_0}) < 0$. Take $\gamma(t) := t^2 t_0^2 u(t t_0 \cdot)$ for $t \in (0, 1]$ and $\gamma(0) := 0$. Then $\gamma \in C([0, 1], H^1(\mathbb{R}^2))$ and $u \in \gamma([0, 1])$. Moreover,

$$\begin{aligned} g'(t) &= t^3 \left[2\|\nabla u\|_2^2 + \frac{1}{t^2} \|u\|_2^2 + \frac{1}{\alpha} \|u\|_2^4 - \frac{(4+\alpha)t^\alpha}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u^2\right) u^2 dx \right. \\ &\quad \left. + \lambda t^{-\frac{3}{2}} \|u\|_2^{5/2} - \frac{2\lambda(r-1)t^{2(r-3)}}{r} \|u\|_r^r - 2 \int_{\mathbb{R}^2} \frac{f(t^2 u)t^2 u - F(t^2 u)}{t^6 u^3} u^3 dx \right], \end{aligned}$$

which implies by (f7) that $t = 1/t_0$ is the unique maximum point of $t \mapsto I_{\alpha,\lambda}(\gamma(t))$. Namely, $\max_{t \in [0, 1]} I_{\alpha,\lambda}(\gamma(t)) = I_{\alpha,\lambda}(u)$. The proof is complete. \square

Lemma 3.7 *Let $\{u_n\} \subset H^1(\mathbb{R}^2)$ be a $(PS)_{c_{\alpha,\lambda}}$ sequence of $I_{\alpha,\lambda}$ for fixed α, λ , then there exists $u_0 \in H^1(\mathbb{R}^2) \setminus \{0\}$ such that $I'_{\alpha,\lambda}(u_0) = 0$.*

Proof In view of Lemma 3.3, we know that $\|u_n\| \leq C$ for some C (independent of n). So there is $u_0 \in H^1(\mathbb{R}^2)$ such that $u_n \rightharpoonup u_0$ weakly in $H^1(\mathbb{R}^2)$. There also exists $B \in \mathbb{R}$ such that

$$\|u_n\|_2^2 \rightarrow B^2, \quad \text{as } n \rightarrow \infty, \tag{3.50}$$

from which we deduce that $J'_{B,\alpha,\lambda}(u_n) \rightarrow 0$ in H^{-1} and $J'_{B,\alpha,\lambda}(u_0) = 0$. In view of Lemma 3.5, for each nontrivial critical point $w^j (j = 1, \dots, k)$ of $J_{B,\alpha,\lambda}$, we have

$$\begin{aligned} \mathcal{B}_{B,\alpha,\lambda}(w^j) &= 2J'_{B,\alpha,\lambda}(w^j)w^j - P_{B,\alpha,\lambda}(w^j) \\ &= 2\|\nabla w^j\|_2^2 + \left(1 + \frac{B^2}{\alpha} + \lambda B^{1/2}\right)\|w^j\|_2^2 - \frac{4+\alpha}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (w^j)^2\right) (w^j)^2 dx \\ &\quad - \left(2 - \frac{2}{r}\right)\lambda \|w^j\|_r^r + 2 \int_{\mathbb{R}^2} [F(w^j) - f(w^j)w^j] dx = 0. \end{aligned} \tag{3.51}$$

Observe from (3.51) that

$$\begin{aligned} &2\|\nabla w^j\|_2^2 + \|w^j\|_2^2 + \frac{1}{\alpha}\|w^j\|_2^4 + \lambda\|w^j\|_2^{5/2} \\ &\leq \frac{4+\alpha}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (w^j)^2\right) (w^j)^2 dx + \left(2 - \frac{2}{r}\right)\lambda \|w^j\|_r^r + 2 \int_{\mathbb{R}^2} [f(w^j)w^j - F(w^j)] dx. \end{aligned} \tag{3.52}$$

From (3.52) and (f₇) we deduce that there exists $t_j \in (0, 1]$ such that

$$\begin{aligned} &2t_j^4\|\nabla w^j\|_2^2 + t_j^2\|w^j\|_2^2 + t_j^4\frac{1}{\alpha}\|w^j\|_2^4 + \lambda t_j^{5/2}\|w^j\|_2^{5/2} \\ &= \frac{4+\alpha}{4\alpha} t_j^{4+\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * |w^j|^2\right) |w^j|^2 dx + \left(2 - \frac{2}{r}\right)\lambda t_j^{2(r-1)}\|w^j\|_r^r \\ &\quad + 2t_j^{-2} \int_{\mathbb{R}^2} [f(t_j^2 w^j)t_j^2 w^j - F(t_j^2 w^j)] dx, \end{aligned} \tag{3.53}$$

which implies $\mathcal{B}_{\alpha,\lambda}(w_{t_j}^j) = 0$, $w_{t_j}^j(x) := t_j^2 w^j(t_j x)$. Then it follows from Lemma 3.6 that there exists $\gamma \in C([0, 1], H^1(\mathbb{R}^2))$ such that $\gamma(0) = 0$, $I_{\alpha,\lambda}(\gamma(1)) < 0$, $w_{t_j}^j \in \gamma([0, 1])$, $0 \notin \gamma((0, 1])$ and

$$\max_{t \in [0, 1]} I_{\alpha,\lambda}(\gamma(t)) = I_{\alpha,\lambda}(w_{t_j}^j).$$

As a result, a direct calculation from (3.53) yields

$$\begin{aligned}
 J_{B,\alpha,\lambda}(w^j) &= J_{B,\alpha,\lambda}(w^j) - \frac{1}{4} \mathcal{B}_{B,\alpha,\lambda}(w^j) \\
 &= \frac{1}{4} \left(1 + \frac{B^2}{\alpha} + \lambda B^{1/2} \right) \|w^j\|_2^2 + \frac{1}{16} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (w^j)^2 \right) (w^j)^2 dx \\
 &\quad + \frac{1}{2} \left(1 - \frac{3}{r} \right) \lambda \|w^j\|_r^r + \frac{1}{2} \int_{\mathbb{R}^2} [f(w^j)w^j - 3F(w^j)] dx \\
 &\geq \frac{1}{4} \|w_{i_j}^j\|_2^2 + \frac{3\lambda}{20} \|w_{i_j}^j\|_2^{5/2} + \frac{1}{16} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * (w_{i_j}^j)^2 \right) (w_{i_j}^j)^2 dx \\
 &\quad + \frac{1}{2} \left(1 - \frac{3}{r} \right) \lambda \|w_{i_j}^j\|_r^r \\
 &\quad + \frac{1}{2} \int_{\mathbb{R}^2} [f(w_{i_j}^j)w_{i_j}^j - 3F(w_{i_j}^j)] dx + \frac{B^2}{4\alpha} \|w^j\|_2^2 + \frac{\lambda B^{1/2}}{10} \|w^j\|_2^2 \\
 &= I_{\alpha,\lambda}(w_{i_j}^j) - \frac{1}{4} \mathcal{B}_{\alpha,\lambda}(w_{i_j}^j) + \left(\frac{B^2}{4\alpha} + \frac{\lambda B^{1/2}}{10} \right) \|w^j\|_2^2 \\
 &\geq c_{\alpha,\lambda} + \left(\frac{B^2}{4\alpha} + \frac{\lambda B^{1/2}}{10} \right) \|w^j\|_2^2.
 \end{aligned} \tag{3.54}$$

Then from Lemma 3.5 and (3.33), we conclude that

$$\begin{aligned}
 c_{\alpha,\lambda} + \frac{B^4}{4\alpha} + \frac{B^{5/2}\lambda}{10} \\
 &= J_{B,\alpha,\lambda}(u_0) + \sum_{j=1}^k J_{B,\alpha,\lambda}(w^j) \\
 &\geq kc_{\alpha,\lambda} + \frac{B^2}{4\alpha} \int_{\mathbb{R}^2} |u_0|^2 dx + \frac{\lambda B^{1/2}}{10} \|u_0\|_2^2 + \left(\frac{B^2}{4\alpha} + \frac{\lambda B^{1/2}}{10} \right) \sum_{j=1}^k \int_{\mathbb{R}^2} |w^j|^2 dx \\
 &\geq kc_{\alpha,\lambda} + \frac{B^4}{4\alpha} + \frac{B^{5/2}\lambda}{10},
 \end{aligned} \tag{3.55}$$

where $w^j \neq 0$ for $j = 1, \dots, k$. Observe that $k > 1$ is impossible.

Thus, $k = 0$, we are done. Then it follows that $J_{B,\alpha,\lambda}(u_0) = I_\alpha(u_0) + \frac{B^4}{4\alpha} + \frac{B^{5/2}\lambda}{10}$ and $u_n \rightarrow u_0$ strongly in $H^1(\mathbb{R}^2)$. Assume $k = 1$ and $u_0 \neq 0$, then the first inequality in (3.55) strictly holds. This yields a contradiction. If $k = 1$ and $u_0 = 0$, then by conclusion (iii) of Lemma 3.5, we get $B = \|w^1\|_2^2$ and $I'_{\alpha,\lambda}(w^1) = 0$ in $H^1(\mathbb{R}^2)$. The proof is complete. \square

4 Proof of Theorem 1.3

In view of Lemma 3.1 and 3.7, there is at least a mountain pass type critical point $u_{\alpha,\lambda}$ of $I_{\alpha,\lambda}$ with $I_{\alpha,\lambda}(u_{\alpha,\lambda}) = c_{\alpha,\lambda}$. That is, $u_{\alpha,\lambda} \in H^1(\mathbb{R}^2)$ is a weak positive solution of equation (3.2).

Choosing a sequence $\{\lambda_n\} \subset (0, 1]$ satisfying $\lambda_n \rightarrow 0^+$, we find a sequence of nontrivial critical points $\{u_{\lambda_n}\}$ (still denoted by $\{u_n\}$) of I_{α,λ_n} with $I_{\alpha,\lambda_n}(u_n) = c_{\alpha,\lambda_n}$. We state the following lemma to ensure that u_n converges strongly to some $u \in H^1(\mathbb{R}^2)$.

Lemma 4.1 For fixed $\alpha \in (0, 1)$, sequence $\{u_n\}$ is bounded in $H^1(\mathbb{R}^2)$.

Proof Multiplying $I_{\alpha,\lambda_n}(u_n)$, $I'_{\alpha,\lambda_n}(u_n)u_n = 0$ and $P_{\alpha,\lambda_n}(u_n) = 0$ by $1, -1/2$ and $1/4$ respectively and adding them up, we get

$$\begin{aligned}
 I_{\alpha,\lambda_n}(u_n) &= \frac{1}{4} \int_{\mathbb{R}^2} u_n^2 dx + \frac{1}{16} Q(u_n) + \frac{3\lambda_n}{20} \|u_n\|_2^{5/2} \\
 &\quad + \frac{1}{2} \int_{\mathbb{R}^2} [f(u_n)u_n - 3F(u_n)] dx + \frac{(r-3)\lambda_n}{2r} \int_{\mathbb{R}^2} |u_n|^r dx,
 \end{aligned}
 \tag{4.1}$$

which implies by Remark 1.2 that $\{u_n\}$ is bounded in $L^2(\mathbb{R}^2)$ uniformly for α, n . Moreover, from (4.1), we have $\int_{\mathbb{R}^2} f(u_n)u_n dx \leq C + 3 \int_{\mathbb{R}^2} F(u_n) dx$. Let $M > \max\{4M_0, t_0\}$, then from (f₁) and (f₅), we conclude that

$$\begin{aligned}
 \int_{\mathbb{R}^2} f(u_n)u_n dx &\leq C + 3 \int_{\mathbb{R}^2} F(u_n) dx \\
 &\leq C + 3 \int_{\{|u_n| \leq M\}} F(u_n) dx + 3 \int_{\{|u_n| \geq M\}} M_0 f(u_n) dx \\
 &\leq C + 3C \int_{\{|u_n| \leq M\}} |u_n|^2 dx + 3 \int_{\{|u_n| \geq M\}} \frac{1}{4} f(u_n)u_n dx,
 \end{aligned}
 \tag{4.2}$$

which implies that $\{f(u_n)u_n\}$ and $\{F(u_n)\}$ are bounded in $L^1(\mathbb{R}^2)$. For $t > 0$, letting $u_{nt}(x) := t^2 u_n(tx)$, we deduce that

$$\begin{aligned}
 &I_{\alpha,\lambda_n}(u_n) - I_{\alpha,\lambda_n}(t^2 u_{nt}) \\
 &= \frac{1-t^4}{2} \|\nabla u_n\|_2^2 + \frac{1-t^2}{2} \int_{\mathbb{R}^2} u_n^2 dx + \lambda_n \frac{2(1-t^{5/2})}{5} \|u_n\|_2^{5/2} \\
 &\quad + \frac{1-t^4}{4\alpha} \|u_n\|_2^4 + \frac{1-t^{4(1+\alpha)}}{4\alpha} Q(u_n) \\
 &\quad + \int_{\mathbb{R}^2} \left[\frac{F(t^2 u_n)}{t^2} - F(u_n) \right] dx - \frac{(1-t^{2(r-1)})\lambda_n}{r} \|u_n\|_r^r \\
 &= \frac{1-t^4}{4} [2I'_{\alpha,\lambda_n}(u_n)u_n - P_{\alpha,\lambda_n}(u_n)] + \left(\frac{3}{20} + \frac{t^4}{4} - \frac{2t^{5/2}}{5} \right) \lambda_n \|u_n\|_2^{5/2} \\
 &\quad + \frac{(1-t^2)^2}{4} \int_{\mathbb{R}^2} u_n^2 dx + \frac{8+\alpha-4t^{4(1+\alpha)}-(4+\alpha)t^4}{16\alpha} Q(u_n) \\
 &\quad + \int_{\mathbb{R}^2} \left[\frac{1-t^4}{2} f(u_n)u_n + \frac{t^4-3}{2} F(u_n) + \frac{1}{t^2} F(t^2 u_n) \right] dx \\
 &\quad + \left[\frac{(r-1)(1-t^4)}{2r} - \frac{(1-t^{2(r-1)})}{r} \right] \lambda_n \|u_n\|_r^r.
 \end{aligned}
 \tag{4.3}$$

We now show that $\{\|\nabla u_n\|_2\}$ is bounded. By contradiction, suppose that $\|\nabla u_n\|_2 \rightarrow \infty$. Take $t_n = (\sqrt{M}/\|\nabla u_n\|_2)^{1/2}$ for some $M > 0$ large, then $t_n \rightarrow 0$. Letting $t = t_n$ in (4.3), since $\{u_n\}$ is bounded in $L^2(\mathbb{R}^2)$, we have

$$\begin{aligned}
 &I_{\alpha,\lambda_n}(u_n) - I_{\alpha,\lambda_n}(t_n^2 u_{nt_n}) \\
 &= \frac{1}{4} \int_{\mathbb{R}^2} u_n^2 dx + \frac{8+\alpha}{16\alpha} Q(u_n) + \int_{\mathbb{R}^2} \left[\frac{1}{2} f(u_n)u_n - \frac{3}{2} F(u_n) + \frac{1}{t_n^2} F(t_n^2 u_n) \right] dx, \\
 &\quad + \frac{r-3}{2r} \lambda_n \|u_n\|_r^r + o(1).
 \end{aligned}
 \tag{4.4}$$

Therefore, it follows from (4.4), Lemma 2.3, the Hardy–Littlewood–Sobolev inequality, and the Gagliardo–Nirenberg inequality that

$$\begin{aligned}
 c_{\alpha,\lambda_n} &\geq I_{\alpha,\lambda_n}(t_n^2 u_n) + \frac{1}{t_n^2} \int_{\mathbb{R}^2} F(t_n^2 u_n) dx + o(1) \\
 &= \frac{t_n^4}{2} \|\nabla u_n\|_2^2 + \frac{t_n^4}{4\alpha} \|u_n\|_2^4 - \frac{t_n^{4+\alpha}}{4\alpha} Q(u_n) \\
 &\quad + \frac{t_n^2}{2} \int_{\mathbb{R}^2} u_n^2 dx + \lambda_n \frac{2t_n^{5/2}}{5} \|u_n\|_2^{5/2} - \frac{t_n^{2r-2}}{r} \lambda_n \|u_n\|_r^r + o(1) \\
 &\geq \frac{t_n^4}{2} \|\nabla u_n\|_2^2 + \frac{t_n^4}{4\alpha} \|u_n\|_2^4 - \frac{t_n^4}{4\alpha} Q(u_n) + o(1) \\
 &\geq \frac{t_n^4}{2} \|\nabla u_n\|_2^2 - \frac{t_n^4}{4} \int \int_{|x-y|\leq 1} \frac{|x-y|^{-\alpha} - 1}{\alpha} u_n^2(x) u_n^2(y) dx dy + o(1) \\
 &\geq \frac{M}{2} - \frac{t_n^4 C}{4} \|u_n\|_2^3 \|\nabla u_n\|_2 + o(1) \\
 &\geq \frac{M}{2} + o(1),
 \end{aligned}
 \tag{4.5}$$

from which we obtain a contradiction by letting $M > 0$ large enough. Hence, $\{u_n\}$ is bounded in $H^1(\mathbb{R}^2)$. □

Remark 4.2 Observe from Lemma 4.1 that $\{u_n\}$ is bounded not only uniformly for n , but also uniformly for α .

Let us assume that $c_{\alpha,\lambda_n} \rightarrow c_\alpha$ as $n \rightarrow \infty$, then $I_{\alpha,\lambda_n}(u_n) \rightarrow c_\alpha$ and $I'_{\alpha,\lambda_n}(u_n) \rightarrow 0$ in H^{-1} . It is easy to see from Lemma 3.4 that $c_\alpha < \frac{2\pi}{\theta_0}$. From Lemma 4.1, we know that $\{u_n\}$ is bounded in $H^1(\mathbb{R}^2)$. Now we take advantage of Lemma 4.1 to get the profile decomposition of $\{u_n\}$. Thus, arguing similarly as in the proof of Lemma 3.5, we have the following lemma.

Lemma 4.3 *Assume that $\{u_n\}$ is a bounded critical point sequence of I_{α,λ_n} with energy level c_α for fixed $\alpha \in (0, 1)$. Then there exist $\tilde{B} \in \mathbb{R}$ and a number $k \in \mathbb{N} \cup \{0\}$, and a finite sequence*

$$(u_0, \tilde{w}^1, \dots, \tilde{w}^k) \subset H^1(\mathbb{R}^2), \quad \tilde{w}^j > 0, \quad \text{for } j = 1, \dots, k \text{ (if } k \geq 1)$$

of critical points for the following functional

$$J_{\tilde{B},\alpha}(u) := \frac{1}{2} \|u\|^2 + \frac{\tilde{B}^2}{2\alpha} \int_{\mathbb{R}^2} |u_n|^2 dx - \frac{1}{4\alpha} \int_{\mathbb{R}^2} \left(\frac{1}{|x|^\alpha} * u^2 \right) u^2 dx - \int_{\mathbb{R}^2} F(u) dx
 \tag{4.6}$$

and k sequences of points $\{\tilde{y}_n^j\} \subset \mathbb{R}^3, 1 \leq j \leq k$, such that

- (i) $|\tilde{y}_n^j| \rightarrow +\infty, |\tilde{y}_n^j - \tilde{y}_n^i| \rightarrow +\infty$ if $i \neq j, n \rightarrow +\infty$,
- (ii) $\|u_n - u_0 - \sum_{j=1}^k \tilde{w}^j(\cdot - \tilde{y}_n^j)\| \rightarrow 0, c_\alpha + \frac{\tilde{B}^4}{4\alpha} = J_{\tilde{B},\alpha}(u_0) + \sum_{j=1}^k J_{\tilde{B},\alpha}(w^j),$
- (iii) $\tilde{B}^2 = \|u_0\|_2^2 + \sum_{j=1}^k \|\tilde{w}^j\|_2^2.$

Otherwise, if $k = 0$, then $u_n \rightarrow u_0$ in $H^1(\mathbb{R}^2)$.

Based on Lemma 4.3, we use the similar arguments as Lemma 3.7 to obtain $u_n \rightarrow u_\alpha$ in $H^1(\mathbb{R}^2)$ as $n \rightarrow \infty$. Moreover, $I'_\alpha(u_\alpha) = 0$ and $I_\alpha(u_\alpha) = c_\alpha$, namely u_α is a nontrivial critical point of I_α . Although we observe from Remark 4.2 that $\|u_\alpha\|$ is uniformly bounded for α , it seems difficult to make use of Moser’s iteration arguments to prove that u_α is bounded in $L^\infty(\mathbb{R}^2)$ uniformly for α , since nonlinearity f is of critical exponential growth in the sense of Trudinger–Moser. More precisely, the following estimate:

$$\sup_{\alpha \in (0, 1)} \|\nabla u_\alpha\|_2^2 < \frac{4\pi}{\theta_0}$$

is not easy to get, so it seems difficult to use the Trudinger–Moser inequality to state a uniform estimate of $f(u_\alpha)$ as $\alpha \rightarrow 0^+$. For fixed $\alpha \in (0, 1)$, arguing as Lemma 3.7 of [36], we can also obtain that there exist C_α, c_α such that

$$u_\alpha(x) \leq C_\alpha \exp(-c_\alpha|x|) \quad \text{for } x \in \mathbb{R}^2.$$

Thus, arguing similarly as Theorem 4.1 in [36], there exists $\alpha_1 \in (0, 1)$ such that for $\alpha \in (0, \alpha_1)$, u_α is radially symmetric up to translation and strictly decreasing in the distance from the symmetry center.

We now state exponential decay estimate of u_α at infinity uniformly for α .

Lemma 4.4 *There exist $R, C > 0$ (independent of α) such that*

$$u_\alpha(x) \leq C \exp\left(-\frac{1}{2}|x|\right) \quad \text{for } |x| \geq R.$$

Proof Since u_α is a positive function, by equation (3.1) and Lemma 2.3 we obtain

$$\begin{aligned} -\Delta u_\alpha + u_\alpha &\leq \int_{|x-y|\leq 1} \frac{|x-y|^{-\alpha}-1}{\alpha} u_\alpha^2(y) dy u_\alpha(x) + f(u_\alpha) \\ &\leq C \int_{|x-y|\leq 1} \frac{u_\alpha^2(y) dy}{|x-y|} u_\alpha(x) + f(u_\alpha). \end{aligned} \tag{4.7}$$

Using the similar arguments as Lemma 2.3 in [36], we have

$$\int_{|x-y|\leq 1} \frac{u_\alpha^2(y)}{|x-y|} dy \rightarrow 0, \quad \text{as } |x| \rightarrow +\infty,$$

uniformly for $\alpha \in (0, \alpha_1)$, which implies that there exists $R_1 > 0$ such that for $|x| \geq R_1$

$$\int_{|x-y|\leq 1} \frac{u_\alpha^2(y)}{|x-y|} dy \leq \frac{1}{4C}. \tag{4.8}$$

By recalling Radial Lemma A.IV in [8], there exists $C > 0$ independent of α such that

$$|u_\alpha(x)| \leq C|x|^{-1} \|u_\alpha\| \leq C|x|^{-1},$$

which implies that

$$\lim_{|x| \rightarrow \infty} |u_\alpha(x)| = 0 \quad \text{uniformly for } \alpha \in (0, \alpha_1).$$

Thus, using assumption (f_1) , we deduce that there exists $R_2 > 0$ such that

$$f(u_\alpha) \leq \frac{1}{4} u_\alpha, \quad |x| \geq R_2. \tag{4.9}$$

Combining (4.7)-(4.9), let $R = \max\{R_1, R_2\}$, then

$$-\Delta u_\alpha + \frac{1}{4}u_\alpha \leq 0, \quad |x| \geq R. \tag{4.10}$$

It follows from (4.10) and a comparison principle, there exists constant $M \geq \frac{C}{R}e^{R/2}$ such that

$$u_\alpha(x) \leq M \exp\left(-\frac{1}{2}|x|\right) \quad \text{for } |x| \geq R.$$

Here R, M are independent of α . The proof is complete. □

Up to a subsequence, we assume

$$\begin{aligned} u_\alpha &\rightharpoonup u_0 \quad \text{in } H^1(\mathbb{R}^2), \\ u_\alpha &\rightarrow u_0 \quad \text{a.e. in } \mathbb{R}^2, \\ u_\alpha &\rightarrow u_0 \quad \text{in } L^s(\mathbb{R}^2) \quad \text{for } s \in (2, +\infty) \end{aligned}$$

as $\alpha \rightarrow 0^+$. For any $\varphi \in C_0^\infty(\mathbb{R}^2)$, we have

$$\begin{aligned} I'_\alpha(u_\alpha)\varphi &= \int_{\mathbb{R}^2} \nabla u_\alpha \nabla \varphi dx + \int_{\mathbb{R}^2} u_\alpha \varphi dx - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|x-y|^{-\alpha} - 1}{\alpha} u_\alpha^2(y) dy u_\alpha(x) \varphi dx \\ &\quad - \int_{\mathbb{R}^2} f(u_\alpha) \varphi dx. \end{aligned} \tag{4.11}$$

Similarly to (3.29), we have

$$\int_{\mathbb{R}^2} f(u_\alpha) \varphi dx \rightarrow \int_{\mathbb{R}^2} f(u_0) \varphi dx, \quad \text{as } \alpha \rightarrow 0^+. \tag{4.12}$$

Then it follows from Lemma 2.3 that for any fixed $\varphi \in C_0^\infty(\mathbb{R}^2)$, we have

$$\left| \frac{|x-y|^{-\alpha} - 1}{\alpha} 1_{|x-y| \leq 1} u_\alpha^2(y) u_\alpha(x) \varphi(x) \right| \leq \left| \frac{1}{|x-y|} u_\alpha^2(y) u_\alpha(x) \varphi(x) \right| := h_\alpha(x, y). \tag{4.13}$$

Since $\{h_\alpha\}$ has a strongly convergent subsequence in $L^1(\mathbb{R}^4)$, we use the Lebesgue dominated convergence theorem to (4.13) to get

$$\begin{aligned} &\int \int_{|x-y| \leq 1} \frac{|x-y|^{-\alpha} - 1}{\alpha} u_\alpha^2(y) dy u_\alpha(x) \varphi(x) dx \rightarrow \\ &\quad - \int \int_{|x-y| \leq 1} \ln|x-y| u_0^2(y) dy u_0(x) \varphi(x) dx. \end{aligned} \tag{4.14}$$

Similarly to the proof of Theorem 1.2 in [36], by Lemma 4.4 and the Lebesgue dominated convergence theorem, one has

$$\begin{aligned} &\int \int_{|x-y| \geq 1} \frac{|x-y|^{-\alpha} - 1}{\alpha} u_\alpha^2(y) dy u_\alpha(x) \varphi(x) dx \rightarrow \\ &\quad - \int \int_{|x-y| \geq 1} \ln|x-y| u_0^2(y) dy u_0(x) \varphi(x) dx. \end{aligned} \tag{4.15}$$

Moreover, by Fatou’s lemma, we have

$$\begin{aligned} & \left| \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y|u_0^2(y)dyu_0^2(x)dx \right| \\ & \leq \liminf_{\alpha \rightarrow 0} \left(\int \int_{|x-y| \leq 1} G_\alpha(x-y)u_\alpha^2(y)dyu_\alpha^2(x)dx - \int \int_{|x-y| \geq 1} G_\alpha(x-y)u_\alpha^2(y)dyu_\alpha^2(x)dx \right). \end{aligned} \tag{4.16}$$

Using Hardy–Littlewood–Sobolev’s inequality and Lemma 2.3, we have

$$\int \int_{|x-y| \leq 1} G_\alpha(x-y)u_\alpha^2(y)dyu_\alpha^2(x)dx < +\infty \tag{4.17}$$

uniformly for α . So by Remark 3.2, we further deduce that

$$\begin{aligned} & \int \int_{|x-y| \geq 1} G_\alpha(x-y)u_\alpha^2(y)dyu_\alpha^2(x)dx \\ & \leq I_\alpha(u_\alpha) + \int \int_{|x-y| \leq 1} G_\alpha(x-y)u_\alpha^2(y)dyu_\alpha^2(x)dx + \int_{\mathbb{R}^2} F(u_\alpha)dx - \frac{1}{2}\|u_\alpha\|^2 \\ & < +\infty \end{aligned} \tag{4.18}$$

uniformly for α . Together (4.16), (4.17) with (4.18), we have

$$\left| \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y|u_0^2(y)dyu_0^2(x)dx \right| < +\infty. \tag{4.19}$$

By virtue of (4.14), (4.15) and (4.19), by taking the limit in (4.11), we have $I'(u_0) = 0$ with $I(u_0) < +\infty$, that is, $u_0 \in H^1(\mathbb{R}^2)$ solves equation (1.4).

We now claim that $u_0 \neq 0$ and $u_\alpha \rightarrow u_0$ in $H^1(\mathbb{R}^2)$. Assume on the contrary that $u_\alpha \rightarrow 0$ in $H^1(\mathbb{R}^2)$, and so $u_\alpha \rightarrow 0$ in $L^s(\mathbb{R}^2)$ for $s \in (2, +\infty)$. Similarly to (3.39), we can obtain $\int_{\mathbb{R}^2} f(u_\alpha)u_\alpha dx = o_\alpha(1)$. So by Lemma 2.3 and Hardy–Littlewood–Sobolev’s inequality, we have

$$\begin{aligned} I'_\alpha(u_\alpha)u_\alpha &= \|u_\alpha\|^2 - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} G_\alpha(x-y)u_\alpha^2(y)u_\alpha^2(x)dx dy - \int_{\mathbb{R}^2} f(u_\alpha)u_\alpha dx \\ &\geq \|u_\alpha\|^2 - \int \int_{|x-y| \leq 1} G_\alpha(x-y)u_\alpha^2(y)u_\alpha^2(x)dx dy + o_\alpha(1) \\ &\geq \|u_\alpha\|^2 - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{|x-y|} u_\alpha^2(y)u_\alpha^2(x)dx dy + o_\alpha(1) \\ &\geq \|u_\alpha\|^2 - \|u_\alpha\|_{\frac{4}{3}}^4 + o_\alpha(1), \end{aligned}$$

which means $u_\alpha \rightarrow 0$ in $H^1(\mathbb{R}^2)$. Then according to Remark 3.2,

$$\begin{aligned} a &\leq I_\alpha(u_\alpha) \\ &= \frac{1}{2}\|u_\alpha\|^2 - \frac{1}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} G_\alpha(x-y)u_\alpha^2(y)u_\alpha^2(x)dx dy - \int_{\mathbb{R}^2} F(u_\alpha)dx \\ &= -\frac{1}{4} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} G_\alpha(x-y)u_\alpha^2(y)u_\alpha^2(x)dx dy + o_\alpha(1) \\ &= o_\alpha(1), \end{aligned}$$

which yields a contradiction and the last identity uses the Lebesgue dominated convergence theorem with Lemma 4.4. Furthermore, similarly to (4.13), (4.15), by Lemma 4.4 and the

Lebesgue dominated convergence theorem, we have

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|x-y|^{-\alpha} - 1}{\alpha} u_\alpha^2(y) u_\alpha^2(x) dy dx \rightarrow - \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \ln|x-y| u_0^2(y) u_0^2(x) dx dy.$$

Using the similar argument as Case 2 of Lemma 3.5, we conclude that $u_\alpha \rightarrow u_0$ in $H^1(\mathbb{R}^2)$ as $\alpha \rightarrow 0^+$. \square

Funding Zhisu Liu is supported by the National Natural Science Foundation of China (No. 12226331), and the Fundamental Research Funds for the Central Universities, China University of Geosciences (Wuhan, No. CUG2106211; CUGST2). The research of Vicențiu D. Rădulescu was supported by a grant of the Romanian Ministry of Research, Innovation and Digitization, CNCS/CCCDI–UEFISCDI, project number PCE 137/2021, within PNCDI III. Jianjun Zhang was supported by the National Natural Science Foundation of China (No. 11871123) and Team Building Project for Graduate Tutors in Chongqing (JDDSTD201802). National Natural Science Foundation of China (11701267).

Data availability No datasets were generated or analysed during the current study.

Code availability Not applicable.

Declarations

Conflict of interest There are no conflict of interest.

Ethical approval Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Adachi, S., Tanaka, K.: Trudinger type inequalities in \mathbb{R}^N and their best exponents. *Proc. Am. Math. Soc.* **128**, 2051–2057 (2000)
2. Adimurthi, Yadava, S.: Multiplicity results for semilinear elliptic equations in bounded domain of \mathbb{R}^2 involving critical exponent. *Ann. Scuola Norm. Super. Pisa-Classe Sci.* **17**, 481–504 (1990)
3. Albuquerque, F., Carvalho, J., Figueiredo, G., Medeiros, E.: On a planar non-autonomous Schrödinger–Poisson system involving exponential critical growth. *Calc. Var. Partial Differ. Equ.* **60**, 30 (2021)
4. Alves, C., Figueiredo, G.: Existence of positive solution for a planar Schrödinger–Poisson system with exponential growth. *J. Math. Phys.* **60**, 011503 (2019)
5. Ambrosetti, A., Ruiz, D.: Multiple bound states for the Schrödinger–Poisson problem. *Commun. Contemp. Math.* **10**, 391–404 (2008)
6. Azzollini, A.: The planar Schrödinger–Poisson system with a positive potential. *Nonlinearity* **34**, 5799–5820 (2021)
7. Benci, V., Fortunato, D.: An eigenvalue problem for the Schrödinger–Maxwell equations. *Topol. Methods Nonlinear Anal.* **11**, 283–293 (1998)
8. Berestycki, H., Lions, P.-L.: Nonlinear scalar field equations. I. Existence of a ground state. *Arch. Ration. Mech. Anal.* **82**, 313–345 (1983)
9. Bernini, F., Mugnai, D.: On a logarithmic Hartree equation. *Adv. Nonlinear Anal.* **9**, 850–865 (2020)
10. Brezis, H., Lieb, E.: A relation between pointwise convergence of functions and convergence of functionals. *Proc. Am. Math. Soc.* **8**, 486–490 (1983)

11. Bucur, C., Cassani, D., Tarsi, C.: Quasilinear logarithmic Choquard equations with exponential growth in R^N . *J. Differ. Equ.* **328**, 261–294 (2022)
12. Cao, D.: Nontrivial solution of semilinear elliptic equation with critical exponent in \mathbb{R}^2 . *Commun. Partial Differ. Equ.* **17**, 407–435 (1992)
13. Cao, D., Dai, W., Zhang, Y.: Existence and symmetry of solutions to 2-D Schrödinger–Newton equations. *Dyn. Partial Differ. Equ.* **18**, 113–156 (2021)
14. Cassani, D., Sani, F., Tarsi, C.: Equivalent Moser type inequalities in \mathbb{R}^2 and the zero mass case. *J. Funct. Anal.* **267**, 4236–4263 (2014)
15. Cassani, D., Tarsi, C.: Schrödinger–Newton equations in dimension two via a Pohozaev–Trudinger log-weighted inequality. *Calc. Var. Partial Differ. Equ.* **60**, 31 (2021)
16. Chen, S., Tang, X.: On the planar Schrödinger–Poisson system with the axially symmetric potential. *J. Differ. Equ.* **268**, 945–976 (2020)
17. Chen, S., Tang, X.: Axially symmetric solutions for the planar Schrödinger–Poisson system with critical exponential growth. *J. Differ. Equ.* **269**, 9144–9174 (2020)
18. Chen, S., Shi, J., Tang, X.: Ground state solutions of Nehari–Pohozaev type for the planar Schrödinger–Poisson system with general nonlinearity. *Discret. Contin. Dyn. Syst. Ser. A* **39**, 5867–5889 (2019)
19. Choquard, P., Stubbe, J., Vuffray, M.: Stationary solutions of the Schrödinger–Newton model—an ODE approach. *Differ. Integral Equ.* **21**, 665–679 (2008)
20. Cingolani, S., Weth, T.: On the planar Schrödinger–Poisson systems. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **33**, 169–197 (2016)
21. Cingolani, S., Jeanjean, L.: Stationary waves with prescribed L^2 -norm for the planar Schrödinger–Poisson system. *SIAM J. Math. Anal.* **51**, 3533–3568 (2019)
22. Dolbeault, J., Frank, R.L., Jeanjean, L.: Logarithmic estimates for mean-field models in dimension two and the Schrödinger–Poisson system. *Comptes Rendus Mathématique* **359**, 1279–1293 (2021)
23. Du, M., Weth, T.: Ground states and high energy solutions of the planar Schrödinger–Poisson system. *Nonlinearity* **30**, 3492–3515 (2017)
24. de Figueiredo, D., Miyagaki, O., Ruf, B.: Elliptic equations in \mathbb{R}^2 with nonlinearities in the critical growth range. *Calc. Var. Partial Differ. Equ.* **3**, 139–153 (1995)
25. Gao, F., Yang, M.: The Brezis–Nirenberg type critical problem for the nonlinear Choquard equation. *Sci. China Math.* **61**, 1219–1242 (2018)
26. He, X.: Multiplicity and concentration of positive solutions for the Schrödinger–Poisson equations. *Z. Angew. Math. Phys.* **62**, 869–889 (2011)
27. Jeanjean, L.: On the existence of bounded Palais–Smale sequence and application to a Landesman–Lazer type problem set on \mathbb{R}^N . *Proc. Roy. Soc. Edinb. Sect. A* **129**, 787–809 (1999)
28. Li, G., Peng, S., Yan, S.: Infinitely many positive solutions for the nonlinear Schrödinger–Poisson system. *Commun. Contemp. Math.* **12**, 1069–1092 (2010)
29. Lieb, E., Loss, M.: *Analysis, Graduate Studies in Mathematics*, vol. 14. American Mathematical Society, Providence, RI (2001)
30. Lions, P.-L.: The concentration compactness principle in the calculus of variations: the locally compact case, Parts 1. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **1**, 109–145 (1984)
31. Lions, P.-L.: The concentration compactness principle in the calculus of variations: The locally compact case. Parts 2. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **2**, 223–283 (1984)
32. Lions, P.-L.: The concentration-compactness principle in the calculus of variations: the limit case 1. *Rev. Iberoam.* **1**, 185–201 (1985)
33. Lions, P.-L.: Solutions of Hartree–Fock equations for Coulomb systems. *Commun. Math. Phys.* **109**, 33–97 (1987)
34. Liu, Z., Zhang, Z., Huang, S.: Existence and nonexistence of positive solutions for a static Schrödinger–Poisson–Slater equation. *J. Differ. Equ.* **266**, 5912–5941 (2019)
35. Liu, Z., Lou, Y., Zhang, J.: A perturbation approach to studying sign-changing solutions of Kirchhoff equations with a general nonlinearity. *Ann. Mat. Pura Appl.* 201 (2022), no. 3, 1229–1255. <https://doi.org/10.1007/s10231-021-01155-w>
36. Liu, Z., Rădulescu, V.D., Tang, C., Zhang, J.: Another look at planar Schrödinger–Newton system. *J. Differ. Equ.* **328**, 65–104 (2022)
37. Moser, J.: A sharp form of an inequality by N. Trudinger. *Indiana Univ. Math. J.* **20**, 1077–1092 (1970–1971)
38. Ruiz, D.: The Schrödinger–Poisson equation under the effect of a nonlinear local term. *J. Funct. Anal.* **237**, 655–674 (2006)
39. Stubbe, J.: Bound states of two-dimensional Schrödinger–Newton equations, e-print, [arXiv:0807.4059](https://arxiv.org/abs/0807.4059)
40. Trudinger, N.S.: On imbeddings into Orlicz spaces and some applications. *J. Math. Mech.* **1**, 473–483 (1967)

41. Wang, Z., Zhou, H.: Positive solution for a nonlinear stationary Schrödinger–Poisson system in \mathbb{R}^3 . *Discret. Contin. Dyn. Syst.* **18**, 809–816 (2007)
42. Wen, L., Chen, S., Rădulescu, V.D.: Axially symmetric solutions of the Schrödinger–Poisson system with zero mass potential in \mathbb{R}^2 . *Appl. Math. Lett.* **104**, 106244, 7 (2020)
43. Willem, M.: *Minimax Theorems*, Progress in Nonlinear Differential Equations and their Applications, vol. 24. Birkhäuser Boston Inc, Boston, MA (1996)
44. Wang, Z.-Q., Zhang, C.: Convergence from power-law to logarithm-law in nonlinear scalar field equations. *Arch. Ration. Mech. Anal.* **231**, 45–61 (2019)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.